

RENORMALIZATION FOR HOLOMORPHIC FIELD THEORIES

BRIAN R. WILLIAMS

ABSTRACT. We introduce the concept of a holomorphic field theory on any complex manifold in the language of the Batalin-Vilkovisky formalism. When the complex dimension is one, this setting agrees with that of chiral conformal field theory. Our main result concerns the behavior of holomorphic theories under renormalization group flow. Namely, we show that holomorphic theories are one-loop finite. We use this to completely characterize holomorphic anomalies in any dimension. Throughout, we compare our approach to holomorphic field theories to more familiar approaches including that of supersymmetric field theories.

CONTENTS

1. Introduction	2
1.1. Acknowledgements	4
2. The definition of a holomorphic field theory	4
2.1. A recollection of the BV-BRST formalism	4
2.2. The definition of a holomorphic field theory	10
2.3. Holomorphically translation invariant theories	22
3. One-loop regularization for theories on \mathbb{C}^d	27
3.1. Homotopy RG flow	30
3.2. Holomorphic gauge fixing	31
3.3. The propagator on \mathbb{C}^d	32
3.4. Trees	34
3.5. A simplification for one-loop weights	35
4. Chiral anomalies in arbitrary dimensions	40
4.1. The quantum master equation	41
4.2. The QME for holomorphic theories	42
Appendix A. Some functional analysis	49
References	51

1. INTRODUCTION

From a mathematical perspective, much of the appeal of quantum field theory is that often theories depend naturally on input geometric data. Furthermore, the usual quantities in quantum field theory such as expectation values and the partition function produce invariants of these underlying geometries. An important and fruitful instance of this is the notion of a *topological field theory*. Mathematically, a topological field theory can be defined on an arbitrary manifold of a fixed dimension. In a precise way, topological theories depend naturally on the smooth structure of the manifold (or smooth structures on associated data such as a bundle). A more complicated class of theories are Riemannian field theories, which, in addition to smooth structures, are sensitive to input metric data. These theories have more refined invariants associated to them, such as the β -function, and are often more relevant to physical examples. In this paper, we study a class of theories that lie between the aforementioned examples. These *holomorphic theories* depend naturally on the complex structure of the underlying space-time.

The idea of studying holomorphic dependence in quantum field theory is certainly not a new one. The most well-known case of this appears in complex dimension one with the notion of a chiral conformal field theory. Here, the holomorphic structure shines most brightly through the *operator product expansion* (OPE) of chiral operators in the theory. This says that the dependence on the product of operators on their relative position is holomorphic, even at the quantum level. These operators combine to form a mathematical object called a vertex algebra. Numerous calculations in conformal field theory reduce to algebraic manipulations at the level of vertex algebras. Furthermore, on arbitrary algebraic curves, the phenomena of operator product expansions has been interpreted mathematically through the pioneering work of Beilinson and Drinfeld on chiral algebras [BD04]. This is arguably one of the greatest successes of mathematics in describing a small, albeit important, class of field theories.

Past dimension one, in complex dimensions two, four and six, an approach to studying special types of holomorphic theories has appeared in the work of Nekrasov and collaborators [Nek96, LMNS97, LMNS96]. There is a holomorphic version of Chern-Simons theory that has appeared in [Pop99, Pop00]. A holomorphic theory of gravity has been proposed in [BCOV94] and studied most recently in the formalism used here in the work of Costello and Costello-Li [CL, Cosa, Cosb]. We will discuss the relationship of our approach to holomorphic field theory to these works throughout this paper.

The goal of this paper is two-fold. In the first part the discussion is fairly formal. After a short recollection of field theory in the Batalin-Vilkovisky formalism, we go on to define the definition of a holomorphic field theory on any complex manifold. We characterize holomorphic deformations of holomorphic theories and provide numerous examples of these theories in the language we set up.

The second part of the paper proceeds to study quantizations of holomorphic field theories defined on \mathbb{C}^d , for any $d \geq 1$. Of course, studying properties of quantization is extremely theory-dependent. Nevertheless, our main result says that when it comes to renormalization, holomorphic theories are generically well-behaved. We show that the renormalization of a holomorphic theory on \mathbb{C}^d is *finite* for quantization at one-loop. A more precise statement is given in Theorem 3.4.

The approach to quantum field theory we use follows Costello’s theory of renormalization and the Batalin-Vilkovisky formalism developed in [Cos11]. In broad strokes, it says that to construct a full quantum field theory it suffices to define the theory at each energy (or length) scale and to ask that these descriptions be compatible as we vary the scale. Concretely, this compatibility is through the *renormalization group (RG) flow* and is encoded by an operator $W(P_{\epsilon < L}, -)$ acting on the space of functionals. The functional $W(P_{\epsilon < L}, -)$ is defined as a sum over weights of graphs which is how Feynman diagrams appear in Costello’s formalism. The infamous infinities of quantum field theory arise due to studying behavior of theories at arbitrarily high energies (or small lengths). In physics this is called the ultra-violet (UV) divergence. Our result can be interpreted by saying that, at one-loop, holomorphic theories have no UV divergences.

Although we do not consider this topic in the present paper, a large collection of examples of holomorphic theories come from familiar physical theories. Namely, holomorphic theories generically appear as minimal *twists* of supersymmetric theories. These are more general than the topological twists considered by Witten in [Wit88]. Any supercharge Q of a supersymmetric theory satisfying $Q^2 = 0$ allows one to construct a “twist”. In some cases, where Clifford multiplication with Q spans all translations such a twist becomes a topological theory (in the weak sense). In any case, however, such a Q defines a “holomorphic twist” [Cos13], which results in the type of holomorphic theories we consider. Regularization in supersymmetric theories, especially gauge theories, is notoriously difficult. Our result implies that after twisting the analytic difficulties become much easier to deal with. Consequently, facets of these theories, such as their anomalies, can be cast in a more algebraic framework. For a recent discussion of holomorphic aspects of twists of supersymmetric theories see [ES].

In no way does this paper tell the complete story of holomorphic field theory. A major future program of the author is to study the behavior of operators for holomorphic field theory, even in the case that the complex manifold is $X = \mathbb{C}^d$. In general, the operators of any quantum field theory form a *factorization algebra* [CG17, CG]. For one-dimensional holomorphic theories, our formalism recovers the theory of chiral and vertex algebras [Wil17, GGW, CG17]. When $d \geq 2$ there is strong evidence that the factorization algebras of holomorphic theories combine to form some higher dimensional vertex algebra structure, where the OPE still varies holomorphically with respect to the relative location of the operators. We will return to this in later publications.

1.1. **Acknowledgements.** The author thanks Owen Gwilliam for comments and suggestions he made on a previous version of this paper that appeared in the author’s thesis. The author would also like to thank Northwestern University, where he received support as a graduate student whilst most of this work took place. In addition, the author enjoyed support as a graduate student research fellow under Award DGE-1324585.

2. THE DEFINITION OF A HOLOMORPHIC FIELD THEORY

The goal of this section is to define the notion of a holomorphic field theory. This is a variant of Costello’s definition of a theory in the Batalin-Vilkovisky formalism, which we will recall at a rapid pace in the first part of this section. In crude summary, to arrive at the definition of a holomorphic field theory we modify the definition of an ordinary BV theory by inserting the word “holomorphic” in front of most objects (bundles, differential operators, etc.). By applying the Dolbeault complex in appropriate locations, we will recover Costello’s definition of a theory, but with a holomorphic flavor, see Table 1.

2.1. **A recollection of the BV-BRST formalism.** In this section we will give an expedient review of the classical Batalin-Vilkovisky formalism. We will also set up the requisite conventions and notations that we will use throughout this paper.

2.1.1. *Classical field theory.* Classical field theory is a formalism for describing a physical system in terms of objects called *fields*. Mathematically, the space of fields is a (most often infinite dimensional) vector space \mathcal{E} . Classical physics is described by the critical locus of a (usually real or complex valued) linear functional on the space of fields

$$(1) \quad S : \mathcal{E} \rightarrow \mathbb{R} \text{ or } \mathbb{C},$$

called the *action functional*. The critical locus is the locus of fields that have zero variation

$$(2) \quad \text{Crit}(S) := \{\varphi \in \mathcal{E} \mid dS(\varphi) = 0\}.$$

A field φ satisfying the equation $dS(\varphi) = 0$ is said to be a *solution to the classical equations of motion*.

Even in the finite dimensional case, if the functional S is not sufficiently well-behaved the critical locus can be still be highly singular. The starting point of the classical Batalin-Vilkovisky formalism is to instead consider the *derived* critical locus. To get a feel for this, we review the finite dimensional situation. Let M be a manifold, which is our ansatz for \mathcal{E} at the moment, and suppose $S : M \rightarrow \mathbb{R}$ is a smooth map. The critical locus is the intersection of the graph of dS in T^*M with the zero section $0 : M \rightarrow T^*M$. Thus, functions on the critical locus are of the form

$$\mathcal{O}(\text{Crit}(S)) = \mathcal{O}(\Gamma(dS)) \otimes_{\mathcal{O}(T^*M)} \mathcal{O}(M).$$

The derived critical locus is a derived space whose dg ring of functions is

$$\mathcal{O}(\text{Crit}^h(S)) = \mathcal{O}(\Gamma(dS)) \otimes_{\mathcal{O}(T^*M)}^{\mathbb{L}} \mathcal{O}(M).$$

We have replaced the strict tensor product with the derived one. Using the Koszul resolution of $\mathcal{O}(M)$ as a $\mathcal{O}(T^*M)$ -module one can write this derived tensor product as a complex of polyvector fields equipped with some differential:

$$\mathcal{O}(\text{Crit}^h(S)) \simeq (\text{PV}^{-*}(M), \iota_{dS}).$$

In cohomological degree $-i$ we have $\text{PV}^{-i}(M) = \Gamma(M, \wedge^i TM)$ and ι_{dS} denotes contraction with the one-form dS (which raises cohomological degree with our regrading convention). With our grading convention we have $\mathcal{O}(T^*[-1]M) = \text{PV}^{-*}(M)$. The space $\mathcal{O}(T^*[-1]M)$ has natural shifted Poisson structure, which takes the form of the familiar Schouten-Nijenhuis bracket of polyvector fields.

The takeaway is that the derived critical locus of a functional $S : M \rightarrow \mathbb{R}$ has the structure of a (-1) -shifted symplectic space. This will be the starting point for our definition of a theory in the BV formalism in the general setting.

In all non-trivial examples the space of fields \mathcal{E} is infinite dimensional and we must be careful with what functionals S we allow. The space of fields we consider will always have a natural topology, and we will choose functionals that are continuous with respect to it. We include a discussion of our convention for infinite dimensional vector spaces including duals and spaces of functionals in the Appendix.

In general, the space of fields of a field theory is equal to the space of smooth sections of a \mathbb{Z} -graded vector bundle $E \rightarrow X$ on a manifold $\mathcal{E} = \Gamma(X, E)$. The \mathbb{Z} -grading is the cohomological, or BRST¹, grading of the theory.

2.1.2. Local functionals. The class of functionals $S : \mathcal{E} \rightarrow \mathbb{R}$ defining the classical theories we consider are required to be *local*, or given by the integral of a Lagrangian density. We define this concept now.

Let D_X denote the sheaf of smooth differential operators on X . If E is any graded vector bundle on X let $\text{Jet}(E)$ denote its bundle of ∞ -jets. This is a smooth vector bundle, albeit infinite rank, on X whose fiber over $y \in X$ can be identified with

$$E_y \times \mathbb{C}[[x_1, \dots, x_n]].$$

Here, $\{x_i\}$ is a formal coordinate near y . This object is given the natural structure of a pro object in the category of vector bundles. We let $J(E)$ denote the associated sheaf of smooth sections. It is well-known that $\text{Jet}(E)$ is equipped with a natural flat connection rendering $J(E)$ with the structure of a smooth D_X -module.

¹Named after Becchi, Rouet, Stora, Tyutin, for which our approach to field theory is greatly influenced by their original mathematical approach to quantization.

In the Appendix we define the algebra of functions $\mathcal{O}(\mathcal{E}(X))$ on the space of global sections $\mathcal{E}(X)$. This is the completed symmetric algebra on the linear dual of $\mathcal{E}(X)$, where the tensor product and dual are interpreted in the appropriate topological sense. Likewise, there is the space of reduced functionals $\mathcal{O}_{red}(\mathcal{E}(X)) = \mathcal{O}(\mathcal{E}(X))/\mathbb{R}$. It is the quotient of all functionals by the constant polynomial functions.

The space $\mathcal{O}_{red}(J(E))$ inherits a natural D_X -module structure from $J(E)$. We refer to $\mathcal{O}_{red}(J(E))$ as the space of *Lagrangians* on the vector bundle E . Every element $F \in \mathcal{O}_{red}(J(E))$ can be expanded as $F = \sum_n F_n$ where each F_n is an element

$$F_n \in \text{Hom}_{C_X^\infty}(J(E)^{\otimes n}, C_X^\infty)_{S_n} \cong \text{PolyDiff}(\mathcal{E}^{\otimes n}, C^\infty(X))_{S_n}$$

where the right-hand side is the space of polydifferential operators. The proof of the isomorphism on the right-hand side can be found in Chapter 5 of [Cos11]. We refer to $\mathcal{O}_{red}(J(E))$ as the (left) D_X -module of *Lagrangians* on the vector bundle E .

A local functional is given by a Lagrangian densities modulo total derivatives. The mathematical definition is the following.

Definition 2.1. Let E be a graded vector bundle on X . Define the sheaf of *local functionals* on X to be

$$\mathcal{O}_{loc}(\mathcal{E}) = \text{Dens}_X \otimes_{D_X} \mathcal{O}_{red}(J(E)),$$

where we use the natural right D_X -module structure on densities.

Note that we always consider local functionals coming from Lagrangians modulo constants. We will not be concerned with local functions associated to constant Lagrangians.

From the expression for functionals in Lemma A.4 we see that integration defines an inclusion of sheaves

$$(3) \quad i : \mathcal{O}_{loc}(\mathcal{E}) \hookrightarrow \mathcal{O}_{red}(\mathcal{E}_c).$$

Often times when we describe a local functional we will write down its value on test compactly supported sections, then check that it is given by integrating a Lagrangian density, which amounts to lifting the functional along i .

2.1.3. *The definition of a classical field theory.* Before giving the definition, we need to recall what the proper notion of a shifted symplectic structure is in the geometric setting that we work in.

Definition 2.2. Let E be a graded vector bundle on X . A *k-shifted symplectic structure* is an isomorphism of graded vector spaces

$$E \cong_\omega E^! [k] = (\text{Dens}_X \otimes E^\vee) [k]$$

that is graded anti-symmetric.

If ω^* is the formal adjoint of the isomorphism $\omega^* : E \cong E^! [k]$, anti-symmetry amounts to the condition $\omega^* = -\omega$. In general, ω does *not* induce a Poisson structure on the space of all functionals $\mathcal{O}(\mathcal{E})$. This is because, as we have seen above, elements of this space are given by distributional sections and hence we cannot pair elements with overlapping support. The symplectic structure does, however, induce a Poisson bracket on *local* functionals.² We will denote the bracket induced by a shifted symplectic structure by $\{-, -\}$.

We are now ready to give the precise definition of a classical field theory.

Definition 2.3 ([CG] Definition 5.4.0.3). *A classical field theory* in the BV formalism on a smooth manifold X is a \mathbb{Z} -graded vector bundle E equipped with a (-1) -shifted symplectic structure together with a local functional $S \in \mathcal{O}_{\text{loc}}(\mathcal{E})$ such that:

- (1) the functional S satisfies the *classical master equation*

$$\{S, S\} = 0;$$

- (2) S is at least quadratic, so we can write it (in a unique way) as

$$S(\varphi) = \omega(\varphi, Q\varphi) + I(\varphi)$$

where Q is a linear differential operator such that $Q^2 = 0$, and $I \in \mathcal{O}_{\text{loc}}(\mathcal{E})$ is at least cubic;

- (3) the complex (\mathcal{E}, Q) is elliptic.

In the physics literature, the operator Q is known as the linearized BRST operator, and $\{S, -\} = Q + \{I, -\}$ is the full BRST operator. Ellipticity of the complex (\mathcal{E}, Q) is a technical requirement that will be very important in our approach to the issue of renormalization in perturbative quantum field theory. The classical master equation is equivalent to

$$QI + \frac{1}{2}\{I, I\} = 0.$$

A *free theory* is a classical theory with $I = 0$ in the notation above. Thus, a free theory is simply an elliptic complex equipped with a (-1) -shifted symplectic pairing where the differential in the elliptic complex is graded skew-self adjoint for the pairing.

Although the space $\mathcal{O}(\mathcal{E})$ does not have a well-defined shifted Poisson bracket induced from the symplectic pairing, the operator $\{S, -\} : \mathcal{O}(\mathcal{E}) \rightarrow \mathcal{O}(\mathcal{E})[1]$ is well-defined since S is local by assumption. By assumption, it is also square zero. The complex of global classical observables of the theory is defined by

$$\text{Obs}_{\mathcal{E}}^{\text{cl}}(X) = (\mathcal{O}(\mathcal{E}(X)), \{S, -\}).$$

This complex is the field theoretic replacement for functions on the derived locus of S from the beginning of this section. Although it does not have a P_0 -structure, there is a

²Note that $\mathcal{O}_{\text{loc}}(\mathcal{E})$ is not a shifted Poisson algebra since there is no natural commutative product.

subspace that does. This is sometimes referred to as the *BRST* complex in the physics literature.

2.1.4. *A description using L_∞ algebras.* There is a completely equivalent way to describe a classical field theory that helps to illuminate the mathematical meaningfulness of the definition given above. The requisite concept we need to introduce is that of a *local Lie algebra* (or local L_∞ algebra).

First, recall that an L_∞ algebra is a modest generalization of a dg Lie algebra where the Jacobi identity is only required to hold up to homotopy. The data of an L_∞ algebra is a graded vector space V with, for each $k \geq 1$, a k -ary bracket

$$\ell_k : V^{\otimes k} \rightarrow V[2 - k]$$

of cohomological degree $2 - k$. These maps are required to satisfy a series of conditions, the first of which says $\ell_1^2 = 0$. The next says that ℓ_2 is a bracket satisfying the Jacobi identity up to a homotopy given by ℓ_3 . For a detailed definition see we refer the reader to [Sta92, Get09].

We now give the definition of a local L_∞ algebra on a manifold X . This has appeared in Chapter 4 of [CG].

Definition 2.4. A *local L_∞ algebra* on X is the following data:

- (i) a \mathbb{Z} -graded vector bundle L on X , whose sheaf of smooth sections we denote \mathcal{L}^{sh} , and
- (ii) for each positive integer n , a polydifferential operator in n inputs

$$\ell_n : \underbrace{\mathcal{L} \times \cdots \times \mathcal{L}}_{n \text{ times}} \rightarrow \mathcal{L}[2 - n]$$

such that the collection $\{\ell_n\}_{n \in \mathbb{N}}$ satisfy the conditions of an L_∞ algebra. In particular, \mathcal{L} is a sheaf of L_∞ algebras.

The simplest example of a local Lie algebra starts with the data of an ordinary Lie algebra \mathfrak{g} . We can then take the constant bundle $\underline{\mathfrak{g}}_X$ with fiber \mathfrak{g} . The Lie bracket on \mathfrak{g} extends to define the structure of a local Lie algebra. In this case, the sheaf of Lie algebras is $C_X^\infty \otimes \mathfrak{g}$. Another important example of a local Lie algebra is given by the Lie algebra of vector fields $\text{Vect}(X)$ on a smooth manifold. The Lie bracket of vector fields is a bidifferential operator on the tangent bundle and this equips the sheaf of sections with the structure of a sheaf of Lie algebras.

Just as in the case of an ordinary graded vector bundle, we can discuss local functionals on a local Lie algebra L . In this case, the L_∞ structure maps give this the structure of a sheaf of complexes, providing a local version of the Chevalley-Eilenberg cochain complex. Indeed, the ∞ -jet bundle JL is an L_∞ algebra object in D_X -modules and so we can define the D_X -module of reduced Chevalley-Eilenberg cochains $C_{\text{Lie,red}}^*(JL)$. Mimicking

the definition above, we arrive at the following local version of Lie algebra cohomology that will come up again and again in this thesis.

Definition 2.5. Let L be a local Lie algebra. The local Chevalley-Eilenberg cochain complex is the sheaf of cochain complexes

$$C_{\text{loc}}^*(\mathcal{L}) = \text{Dens}_X \otimes_{D_X} C_{\text{Lie,red}}^*(L).$$

We denote the global sections by $C_{\text{loc}}^*(\mathcal{L}(X))$.

The *local cohomology* of a local Lie algebra is the cohomology of the local CE complex, which we will denote $H_{\text{loc}}^*(\mathcal{L}(X))$.

Remark 2.6. We have already remarked that for a graded vector bundle E there is an embedding $\mathcal{O}_{\text{loc}}(\mathcal{E}) \hookrightarrow \mathcal{O}_{\text{red}}(\mathcal{E})$. This translates to an embedding of sheaves of cochain complexes $C_{\text{loc}}^*(\mathcal{L}) \hookrightarrow C_{\text{Lie,red}}^*(\mathcal{L}_c)$ for any local Lie algebra \mathcal{L} . In the case of vector fields, there is a related cochain complex that has been studied extensively in the context of characteristic classes of foliations [Fuk86, Gui73, Los98, BR73]. Suppose, for simplicity, that X is a compact smooth manifold. If $\text{Vect}(X)$ is the Lie algebra of vector fields on X then the (reduced) *diagonal cochain complex* is the subcomplex

$$C_{\Delta,\text{red}}^*(\text{Vect}(X)) \subset C_{\text{Lie,red}}^*(\text{Vect}(X))$$

consisting of cochains $\varphi : \text{Vect}(X)^{\otimes k} \rightarrow \mathbb{C}$ satisfying $\varphi(X_1, \dots, X_k) = 0$ if $\bigcap_{i=1}^k \text{Supp}(X_i) = \emptyset$. That is, the cocycle vanishes unless all of the supports of the inputs overlap nontrivially. The inclusion of the local cochain complex $C_{\text{loc}}^*(\text{Vect}(X)) \subset C_{\text{Lie,red}}^*(\text{Vect}(X))$ factors through this subcomplex to give a sequence of inclusions

$$C_{\text{loc}}^*(\text{Vect}(X)) \hookrightarrow C_{\Delta,\text{red}}^*(\text{Vect}(X)) \hookrightarrow C_{\text{Lie,red}}^*(\text{Vect}(X)).$$

This is because the cochain of $\text{Vect}(X)$ defined from a local cochain involves the integral of local operators applied to the inputs.

It turns out that the definition of a classical field theory can be repackaged in terms of certain structures on a local L_∞ algebra. The first piece of data we need to transport to the L_∞ side is that of a symplectic pairing. The underlying data of a local L_∞ algebra L is a graded vector bundle. In Definition 2.2 we have already defined a k -shifted symplectic pairing. On the local Lie algebra sign, we ask for $k = -3$ shifted symplectic structures that are also invariant for the L_∞ structure maps.

Also, an important part of a classical field theory is ellipticity. We say a local L_∞ algebra is *elliptic* if the complex $(\mathcal{L}, d = \ell_1)$ is an elliptic complex.

Proposition 2.7 ([CG] Proposition 5.4.0.2). *The following structures are equivalent:*

- (1) a classical field theory in the BV formalism (\mathcal{E}, ω, S) ;
- (2) an elliptic local Lie algebra structure on $L = E[1]$ equipped with a (-3) -shifted symplectic pairing.

Proof. (Sketch) The underlying graded vector bundle of the space of fields \mathcal{E} is E and we obtain the bundle underlying the local L_∞ algebra by shifting this down $L = E[1]$. The (-1) -shifted symplectic structure on E transports to a (-3) -shifted on L . The L_∞ structure maps for L come from the Taylor components of the action functional S . The exterior derivative of S is a section

$$dS \in C_{\text{loc}}^*(\mathcal{L}, \mathcal{L}^![-1]),$$

where on the right-hand side we have zero differential. The Taylor components are of the form $(dS)_n : \mathcal{L}^{\otimes n} \rightarrow \mathcal{L}^![-1]$. Using the shifted symplectic pairing we can identify these Taylor components with maps $(dS)_n : \mathcal{L}^{\otimes n} \rightarrow \mathcal{L}[2]$. Thus, dS can be viewed as a section of $C_{\text{loc}}^*(\mathcal{L}, \mathcal{L}[2])$. This is precisely the space controlling deformations of \mathcal{L} as a local Lie algebra. One checks immediately that the classical master equation is equivalent to the fact that dS is a derivation, hence it determines the structure of a local Lie algebra. The first Taylor component ℓ_1 is precisely the operator Q before, so ellipticity of (\mathcal{E}, Q) is equivalent to ellipticity of (\mathcal{L}, ℓ_1) . \square

2.2. The definition of a holomorphic field theory. In this section we fix a complex manifold X of complex dimension d . We start with the definition of a *free* holomorphic field theory on X , from there we will go on to describe how to incorporate interactions.

2.2.1. Free holomorphic theories. The essential information that governs a classical field theory are its equations of motion. For a free theory, the equations of motion are linear in the space of fields. At least classically, the setting of free theories can essentially be reduced to the study linear partial differential equations.

First, we must come to terms with the fields of a holomorphic theory. Just as in the case of an ordinary field theory, they will arise as sections of some \mathbb{Z} -graded vector bundle on X . The \mathbb{Z} -grading plays the same role as in the usual setting, it counts the BRST, or ghost, degree. We will also refer to this as the cohomological degree. For a *holomorphic theory* the crucial step is that we impose that this graded vector bundle be holomorphic. By a holomorphic \mathbb{Z} -graded vector bundle we mean a \mathbb{Z} -graded vector bundle $V^\bullet = \bigoplus_i V^i[-i]$ (which we will usually abbreviate simply as V) such that each graded piece V^i is a holomorphic vector bundle (here V^i is in cohomological degree $+i$). Thus, in order to define a holomorphic field theory on a complex manifold X we start with the data:

- (1) a \mathbb{Z} -graded holomorphic vector bundle $V^\bullet = \bigoplus_i V^i[-i]$ on X , so that the finite dimensional holomorphic vector bundle V^i is in cohomological degree i .

Remark 2.8. For supersymmetric theories it may be desirable to include an additional $\mathbb{Z}/2$, or fermionic, grading into the data of the space of fields, but we do not consider that here.

A free classical theory is made up of a space of fields as above together with the data of a linearized BRST differential Q^{BRST} and a shifted symplectic pairing of cohomological degree -1 . Ordinarily, the BRST operator is simply a differential operator on the underlying vector bundle defining the fields. For the class of theories we are considering, we require this operator be holomorphic. For completeness, we briefly recall this notion.

Suppose that E and F are two holomorphic vector bundles on X . Note that the Hom-bundle $\text{Hom}(E, F)$ inherits a natural holomorphic structure. By definition, a *holomorphic differential operator of order m* is a linear map

$$D : \Gamma^{hol}(X; E) \rightarrow \Gamma^{hol}(X; F)$$

such that, with respect to a holomorphic coordinate chart $\{z_i\}$ on X , D can be written as

$$(4) \quad D|_{\{z_i\}} = \sum_{|I| \leq m} a_I(z) \frac{\partial^{|I|}}{\partial z_I}$$

where $a_I(z)$ is a local holomorphic section of $\text{Hom}(E, F)$. Here, the sum is over all multi-indices $I = (i_1, \dots, i_d)$ and

$$\frac{\partial^{|I|}}{\partial z_I} := \prod_{k=1}^d \frac{\partial^{i_k}}{\partial z_k^{i_k}}.$$

The length of the multi-index I is defined by $|I| := i_1 + \dots + i_d$.

Example 2.9. The most basic example of a holomorphic differential operator is the ∂ operator for the trivial vector bundle. For each $1 \leq \ell \leq d = \dim_{\mathbb{C}}(X)$, it is a holomorphic differential operator from $E = \wedge^{\ell} T^{1,0*} X$ to $F = \wedge^{\ell+1} T^{1,0*} X$ which on sections is

$$\partial : \Omega^{\ell, hol}(X) \rightarrow \Omega^{\ell+1, hol}(X).$$

Locally, of course, it has the form

$$\partial = \sum_{i=1}^d (dz_i \wedge (-)) \frac{\partial}{\partial z_i},$$

where $dz_i \wedge (-)$ is the vector bundle homomorphism $\wedge^{\ell} T^{1,0*} X \rightarrow \wedge^{\ell+1} T^{1,0*} X$ sending $\alpha \mapsto dz_i \wedge \alpha$.

The next piece of data we fix is:

- (2) a square-zero holomorphic differential operator

$$Q^{hol} : \mathcal{V}^{hol} \rightarrow \mathcal{V}^{hol}$$

of cohomological degree $+1$. Here \mathcal{V}^{hol} denotes the holomorphic sections of V .

Finally, to define a free theory we need the data of a shifted symplectic pairing. For reasons to become clear in a moment, we must choose this pairing to have a strange cohomological degree. The last piece of data we fix is:

(3) an invertible bundle map

$$(-, -)_V : V \otimes V \rightarrow K_X[d - 1]$$

Here, K_X is the canonical bundle on X .

The definition of the fields of an ordinary field theory are the *smooth* sections of the vector bundle V . In our situation this is a silly thing to do since we lose all of the data of the complex structure we used to define the objects above. The more natural thing to do is to take the *holomorphic* sections of the vector bundle V . By construction, the operator Q^{hol} and the pairing $(-, -)_V$ are defined on holomorphic sections, so on the surface this seems reasonable. The technical caveat that the sheaf of holomorphic sections does not satisfy certain conditions necessary to study renormalization and observables in our approach to QFT. For more details on this see Remark 2.12. The solution to this problem is to take a natural resolution of holomorphic sections in order to relate to the usual definition of a classical BV theory.

Given any holomorphic vector bundle V we can define its *Dolbeault complex* $\Omega^{0,*}(X, V)$ with its Dolbeault operator

$$\bar{\partial} : \Omega^{0,p}(X, V) \rightarrow \Omega^{0,p+1}(X, V).$$

Here, $\Omega^{0,p}(X, V)$ denotes smooth sections of the vector bundle $\wedge^p(T^{0,1})^\vee X \otimes V$. For any $U \subset X$ open subset, the complex $\Omega^{0,*}(U, V)$ is defined. In this way, we obtain a natural sheaf of complexes on X , that we denote by $\Omega_X^{0,*}(V)$. The fundamental property of the Dolbeault complex is that by Dolbeault's Theorem it provides a resolution for the sheaf of holomorphic sections:

$$\mathcal{V}^{hol} \rightarrow \Omega_X^0(V) \xrightarrow{\bar{\partial}} \Omega_X^{0,1}(V) \xrightarrow{\bar{\partial}} \dots$$

We now take a graded holomorphic vector bundle $V = V^\bullet$ as above, equipped with the differential operator Q^{hol} . The Dolbeault resolution $\Omega^{0,*}(X, V^\bullet)$ is now equipped with two differentials Q^{hol} and $\bar{\partial}$. The complex of fields is the totalization of this complex:

$$\mathcal{E}_V = \text{Tot} \left(\Omega^{0,*}(X, V), \bar{\partial}, Q^{hol} \right) = \left(\Omega^{0,*}(X, V), \bar{\partial} + Q^{hol} \right).$$

The operator $\bar{\partial} + Q^{hol}$ will be the linearized BRST operator of our theory. By assumption, we have $[\bar{\partial}, Q^{hol}]$ so that $(\bar{\partial} + Q^{hol})^2 = 0$ and hence the fields still define a complex.

By construction, \mathcal{E}_V has the natural structure of a sheaf of complexes. When we want to consider global sections over X we use the notation $\mathcal{E}_V(X)$. There is similarly a cosheaf of compactly supported sections $\mathcal{E}_{V,c}$ whose underlying graded is the compactly supported Dolbeault forms $\Omega_c^{0,*}(X, V)$.

The pairing $(-, -)_V$ defines a pairing on \mathcal{E}_V as follows. The thing to observe here is that $(-, -)_V$ extends to the Dolbeault complex in a natural way: we simply combine the

wedge product of forms with the pairing on V . We obtain the following composition.

$$\begin{array}{ccc} \mathcal{E}_{V,c} \otimes \mathcal{E}_{V,c} & \xrightarrow{(-,-)_V} & \Omega_c^{0,*}(X, K_X)[d-1] \\ & \searrow \omega_V & \downarrow \int_X \\ & & \mathbb{C}[-1]. \end{array}$$

The top Dolbeault forms with values in the canonical bundle K_X are precisely the top forms on the smooth manifold X , and we use the integration map $\int_X : \Omega_c^{d,d}(X) \rightarrow \mathbb{C}$. We note that integration is of cohomological degree d , as exhibited in the diagram.

We arrive at the following definition.

Definition/Lemma 1. A free holomorphic theory on a complex manifold X is the data $(V, Q^{hol}, (-, -)_V)$ as in (1), (2), (3) above such that Q^{hol} is a square zero elliptic differential operator that is graded skew self-adjoint for the pairing $(-, -)_V$. The triple $(\mathcal{E}_V, Q_V = \bar{\partial} + Q^{hol}, \omega_V)$ defines a free BV theory in the usual sense.

The usual prescription for writing down the associated action functional holds in this case. If $\varphi \in \Omega^{0,*}(X, V)$ denotes a field the action is

$$S(\varphi) = \int_X \left(\varphi, (\bar{\partial} + Q^{hol})\varphi \right)_V.$$

We arrive at an example, which is a higher dimensional version of a familiar chiral CFT.

Example 2.10. The free $\beta\gamma$ system. Suppose that

$$V = \underline{\mathbb{C}} \oplus K_X[d-1].$$

Let $(-, -)_V$ be the pairing

$$(\underline{\mathbb{C}} \oplus K_X) \otimes (\underline{\mathbb{C}} \oplus K_X) \rightarrow K_X \oplus K_X \rightarrow K_X$$

sending $(\lambda, \mu) \otimes (\lambda', \mu') \mapsto (\lambda\mu', \lambda'\mu) \mapsto \lambda\mu' + \lambda'\mu$. In this example we set $Q^{hol} = 0$. One immediately checks that this is a holomorphic free theory as above. The space of fields can be written as

$$\mathcal{E}_V = \Omega^{0,*}(X) \oplus \Omega^{d,*}(X)[d-1].$$

We write $\gamma \in \Omega^{0,*}(X)$ for a field in the first component, and $\beta \in \Omega^{d,*}(X)[d-1]$ for a field in the second component. The action functional reads

$$S(\gamma + \beta, \gamma' + \beta') = \int_X \beta \wedge \bar{\partial}\gamma' + \beta' \wedge \bar{\partial}\gamma.$$

When $d = 1$ this reduces to the ordinary chiral $\beta\gamma$ system from conformal field theory. The $\beta\gamma$ system is a bosonic version of the ghost bc system that appears in the quantization of the bosonic string, see Chapter 6 of [Pol98]. For instance, we will see how this theory is the starting block for constructing general holomorphic σ -models.

Example 2.11. Coefficients in a bundle. There are many variants of the $\beta\gamma$ system that we can consider. For instance, if E is any (graded) holomorphic vector bundle on X we can take

$$V = E \oplus K_{\mathbb{C}^d} \otimes E^\vee$$

where E^\vee is the linear dual bundle. The pairing is constructed as in the case above where we also use the evaluation pairing between E and E^\vee . In this case, the fields are $\gamma \in \Omega^{0,*}(X, E)$ and $\beta \in \Omega^{d,*}(X, E^\vee)[d-1]$. The action functional is simply

$$S(\gamma, \beta) = \int \text{ev}_E(\beta \wedge \bar{\partial}\gamma).$$

Here, ev_E stands for the evaluation pairing between sections of E and sections of the dual E^\vee . When E is a tensor bundle of type (r, s) this theory is a bosonic version of the bc ghost system of spin (r, s) . For a general bundle E we will refer to it as the $\beta\gamma$ system with coefficients in the bundle E .

Remark 2.12. We will only work with a holomorphic theory prescribed by the data $(V, (-, -)_V, Q^{hol})$ through its associated BV theory. One might propose a definition of a BV theory in the analytic category based off of holomorphic sections of holomorphic vector bundles. There are numerous technical reason why this approach fails in our approach to QFT. In particular, the sheaf of holomorphic sections of a holomorphic bundle is not a fine sheaf, and there do not exist partitions of unity in general. In addition, there is no holomorphic analog of compactly supported smooth functions. Compact support is crucial when consider locality in field theory. For instance, the main result of [CG] is that the observables of any QFT form a factorization algebra, which is heavily on the existence of sections with compact support. A dual approach to higher dimensional versions of chiral algebras, in the style of Beilinson and Drinfeld, has been proposed in [FG12]. This formulation makes sense in the algebraic or analytic category and should be connected to our anticipated version of chiral algebras as observables of a holomorphic field theory.

2.2.2. Interacting holomorphic theories. We proceed to define what an interacting holomorphic theory is. A general interacting field theory with space of fields \mathcal{E} is prescribed by a functional

$$S : \mathcal{E} \rightarrow \mathbb{C}$$

that satisfies the *classical master equation*. The key technical condition is that this functional must, in addition, be *local*. For a precise definition see Chapter 5 of [CG].

Since X is a complex manifold, it makes sense to consider the sheaf of holomorphic differential operators that we denote by D_X^{hol} . If V is a holomorphic vector bundle we define the bundle of holomorphic ∞ -jets $\text{Jet}^{hol}(V)$ as follows [GG80, CW04]. This is a pro-vector bundle that is holomorphic in a natural way. The fibers of this infinite rank bundle $\text{Jet}^{hol}(V)$ are isomorphic to

$$\text{Jet}^{hol}(V)|_w = V_w \otimes \mathbb{C}[[z_1, \dots, z_d]],$$

where $w \in X$ and where $\{z_i\}$ is the choice of a holomorphic formal coordinate near w . We denote by $J^{hol}V$ the sheaf of holomorphic sections of this jet bundle. The sheaf $J^{hol}V$ has the structure of a D_X^{hol} -module, that is, it is equipped with a holomorphic flat connection ∇^{hol} . This situation is completely analogous to the smooth case. Locally, the holomorphic flat connection on $\text{Jet}^{hol}(V)$ is of the form

$$\nabla^{hol}|_w = \sum_{i=1}^d dw_i \left(\frac{\partial}{\partial w_i} - \frac{\partial}{\partial z_i} \right),$$

where $\{w_i\}$ is the local coordinate on X near w and z_i is the fiber coordinate labeling the holomorphic jet expansion.

One natural appearance of the bundle of holomorphic jets is in providing an explicit description of holomorphic differential operators. The statement in the smooth category is simply that a differential operator between vector bundles is equivalent to the data of a map of D -modules between the associated ∞ -jet bundles. In a completely analogous way, holomorphic differential operators are the same as bundle maps between the associated holomorphic jet bundles. A similar result holds for *polydifferential* operators, which we also state.

Lemma 2.13. *Suppose V, W are holomorphic vector bundles with spaces of holomorphic sections given by $\mathcal{V}^{hol}, \mathcal{W}^{hol}$ respectively. There is an isomorphism of sheaves on X*

$$\text{Diff}^{hol}(\mathcal{V}^{hol}, \mathcal{W}^{hol}) \cong \text{Hom}_{D_X^{hol}}(J^{hol}(V), J^{hol}(W)).$$

Similarly, if V_1, \dots, V_n, W are holomorphic bundles on X , there is an isomorphism

$$\text{PolyDiff}^{hol}(\mathcal{V}_1^{hol} \times \dots \times \mathcal{V}_n^{hol}, \mathcal{W}^{hol}) \cong \text{Hom}(J^{hol}(V_1) \otimes \dots \otimes J^{hol}(V_n), W).$$

In both cases, the right-hand side denotes the space of homomorphisms of holomorphic D -modules that are compatible with the adic topology on jets.

We will utilize this interpretation of holomorphic jet bundles momentarily.

In ordinary field theory, local functionals are defined as integrals of Lagrangian densities. By definition, a Lagrangian density is a density valued functional on the fields that only depends on the fields through it's partial derivatives. In the holomorphic setting we have the following definition.

Definition 2.14. Let V be a vector bundle. The space of *holomorphic Lagrangian densities* on V is

$$\text{Lag}^{hol}(V) = \Omega_X^{d,hol} \otimes_{\mathcal{O}_X^{hol}} \left(\prod_{n>0} \text{Hom}_{\mathcal{O}_X^{hol}}(J^{hol}(V)^{\otimes n}, \mathcal{O}_X^{hol})_{S_n} \right).$$

The hom space inside the parentheses denotes maps of holomorphic vector bundles respecting the natural filtration on jets. That is, we require the bundle maps to be continuous with respect to the natural adic topology. We also take coinvariants for the symmetric group S_n .

Note that we take the product over $n > 0$. We do not want to consider Lagrangians that are constant in the fields.

Equivalently, a holomorphic Lagrangian density is of the form $\omega \otimes F$ where ω is a top holomorphic form and F is a functional $F = \sum_k F_k$ where, for each k , the multilinear map

$$F_k : \mathcal{V}^{hol} \times \dots \times \mathcal{V}^{hol} \rightarrow \mathcal{O}_X^{hol}$$

depends only on the holomorphic ∞ -jet of sections of V .

Next, suppose that V is part of the data of a free holomorphic theory $(V, Q^{hol}, (-, -)_V)$. The pairing $(-, -)_V$ endows the space of holomorphic Lagrangians with a sort of bracket that we now construct.

Suppose $\omega \otimes F, \omega' \otimes F' \in \text{Lag}^{hol}(V)$. For simplicity, we assume F, F' are of homogeneous symmetric degree k, k' respectively. Consider the natural map

$$(5) \quad \left(\Omega_X^{d,hol} \otimes \text{Hom}(J^{hol}(V)^{\otimes k}, \mathcal{O}_X^{hol}) \right) \otimes \left(\Omega_X^{d,hol} \otimes \text{Hom}(J^{hol}(V)^{\otimes k'}, \mathcal{O}_X^{hol}) \right) \\ \downarrow \\ \left(\Omega_X^{d,hol} \otimes \Omega_X^{d,hol} \right) \otimes \text{Hom}(J^{hol}(V)^{\otimes (k+k')}, \mathcal{O}_X^{hol})$$

for which the product $(\omega \otimes F) \otimes (\omega' \otimes F')$ maps to $(\omega \otimes \omega') \otimes (F \otimes F')$. The bundle map $(-, -)_V : V \otimes V \rightarrow K_X[d-1]$ is invertible, hence it determines a section $(-, -)_V^{-1}$ of $V \otimes V \otimes K_X^\vee$ of cohomological degree $d-1$, where K_X^\vee is the dual bundle to K_X . Sections of K_X^\vee will be denoted PV_X^d (the PV stands for polyvector fields). The element $(-, -)_V^{-1}$ determines a map

$$(6) \quad \text{Hom}(J^{hol}(V)^{\otimes (k+k')}, \mathcal{O}_X^{hol}) \xrightarrow{(-, -)_V^{-1}} \text{Hom}(J^{hol}(V)^{\otimes (k+k'-2)}, \text{PV}_X^d).$$

Explicitly, this sends the element $F \otimes F'$ to the functional

$$\varphi_1 \otimes \dots \otimes \varphi_{k+k'-2} \mapsto (F \otimes F' \otimes \text{id}_{K_X^\vee})(\varphi_1 \otimes \dots \otimes \varphi_{k+k'-2} \otimes (-, -)_V^{-1}) \in \text{PV}_X^d.$$

Finally, using the natural pairing between top polyvector fields and top holomorphic forms, we obtain a map

$$(7) \quad \left(\Omega_X^{d,hol} \otimes \Omega_X^{d,hol} \right) \otimes \text{Hom}(J^{hol}(V)^{\otimes (k+k'-2)}, \text{PV}_X^d) \rightarrow \Omega_X^{d,hol} \otimes \text{Hom}(J^{hol}(V)^{\otimes (k+k')}, \mathcal{O}_X^{hol}).$$

Upon symmetrizing the inputs, we see that the image of $(\omega \otimes F) \otimes (\omega' \otimes F')$ under the composition of the maps (5), (6), and (7) determines an element $\{F, F'\}^{hol} \in \text{Lag}^{hol}(V)$. We extend to non-homogenous functionals in the obvious way.

In conclusion, we have produced a bilinear map

$$\{-, -\}^{hol} : \text{Lag}^{hol}(V) \times \text{Lag}^{hol}(V) \rightarrow \text{Lag}^{hol}(V)[d-1].$$

Note that this bracket is of cohomological degree $-d+1$ and lowers the polynomial degree by two.

We can now state the definition of a classical holomorphic theory. Note that top holomorphic forms have a natural action by the Lie algebra of holomorphic vector fields \mathcal{T}_X^{hol} via Lie derivative. This induces an action of holomorphic vector fields on the space of holomorphic Lagrangians.

Definition 2.15. A *classical holomorphic theory* on a complex manifold X is the data of a free holomorphic theory $(V, Q^{hol}, (-, -)_V)$ plus a holomorphic Lagrangian

$$I^{hol} \in \Omega_X^{d, hol} \otimes_{\mathcal{O}_X^{hol}} \prod_{n \geq 3} \text{Hom}(J^{hol}(V)^{\otimes n}, \mathcal{O}_X^{hol})_{S_n} \subset \text{Lag}^{hol}(V)$$

of cohomological degree d such that the following holomorphic Lagrangian

$$Q^{hol} I^{hol} + \frac{1}{2} \{I^{hol}, I^{hol}\}^{hol}$$

is in the image of a some holomorphic vector field. That is, there exists some $\xi \in \mathcal{T}_X^{hol}$ and $F \in \text{Lag}^{hol}(V)$ such that $Q^{hol} I^{hol} + \frac{1}{2} \{I^{hol}, I^{hol}\}^{hol} = L_\xi F$.

Remark 2.16. Note that in the definition we require that the functional I^{hol} be at least cubic. For brevity, we will denote the subspace $\prod_{n \geq 3} \text{Hom}(\text{Jet}^{hol}(V)^{\otimes n}, K_X)_{S_n}$ by $\text{Lag}^{hol,+}(V)$.

There is an alternative way to understand the condition on the functional I^{hol} to define a classical holomorphic theory. To define it, we introduce the notion of a holomorphic local functional. A holomorphic local functional is, by definition, a holomorphic Lagrangian defined up to a total holomorphic derivative. Precisely, we have the following definition.

Definition 2.17. The sheaf of holomorphic local functionals is defined to be the quotient

$$\mathcal{O}_{loc}^{hol}(V) := \text{Lag}^{hol}(V) / \mathcal{T}_X^{hol} \cdot \text{Lag}^{hol}(V),$$

where $\mathcal{T}_X \cdot \text{Lag}^{hol}(V)$ denotes the subspace of holomorphic Lagrangians that are in the image of the Lie derivative by some holomorphic vector field.

Equivalently, we may express this quotient using holomorphic D -modules in the following way. The left D_X^{hol} -module structure on $J^{hol}(V)$ carries over to a left D_X^{hol} -module structure on the product

$$\prod_{n > 0} \text{Hom}_{\mathcal{O}_X^{hol}}(J^{hol}(V)^{\otimes n}, \mathcal{O}_X^{hol}).$$

Moreover, we have already mentioned that there is the natural structure of a right D_X^{hol} -module structure on $\Omega_X^{d, hol}$.

Lemma 2.18. *There is an isomorphism*

$$\mathcal{O}_{loc}^{hol}(V) = \Omega_X^{d, hol} \otimes_{D_X^{hol}} \prod_{n > 0} \text{Hom}_{\mathcal{O}_X^{hol}}(J^{hol}(V)^{\otimes n}, \mathcal{O}_X^{hol}).$$

Compare this to the definition of ordinary local functionals that we recalled in Definition 2.1.

Next, we have the following observation.

Lemma 2.19. *Both the bracket $\{-, -\}^{hol}$ and the differential Q^{hol} on holomorphic Lagrangians descends to a bracket and differential on $\mathcal{O}_{loc}^{hol}(V)$ (that we denote by the same names). This yields the structure of a dg Lie algebra*

$$(\mathcal{O}_{loc}^{hol}(V)[d-1], Q^{hol}, \{-, -\}^{hol}).$$

An immediate corollary of this lemma is that the condition for a holomorphic Lagrangian I^{hol} to define a classical theory in Definition 2.17 is equivalent to the condition that I^{hol} is a Maurer-Cartan element in $\mathcal{O}_{loc}^{hol}(V)[d-1]$.

As in the free case, we proceed to verify that a holomorphic theory defines an interacting classical BV theory in the sense of Definition 2.3.

The underlying space of fields, as we have already seen in the free case, is $\mathcal{E}_V = \Omega^{0,*}(X, V)$. We show how to extend a holomorphic Lagrangian to a functional on this Dolbeault complex.

Recall, a holomorphic Lagrangian can be written as $I^{hol} = \sum_k I_k^{hol}$ where $I_k^{hol} = \omega \otimes F_k$ for $\omega \in \Omega^{d,hol}$ and $F_k : \mathcal{V}^{hol} \times \dots \times \mathcal{V}^{hol} \rightarrow \mathcal{O}^{hol}$ is of the form

$$F_k(\varphi_1, \dots, \varphi_k) = \sum_{i_1, \dots, i_k} D_{i_1}(\varphi_1) \cdots D_{i_k}(\varphi_k) \in \mathcal{O}_X^{hol}.$$

Here, $\varphi_i \in \mathcal{V}^{hol}$ is a holomorphic section, and each D_{i_j} is a holomorphic differential operator $D_{i_j} : \mathcal{V}^{hol} \rightarrow \mathcal{O}^{hol}$.

In general, suppose V, W are holomorphic vector bundles. Every holomorphic differential operator $D : \mathcal{V} \rightarrow \mathcal{W}$ extends to a *smooth* differential operator on the associated Dolbeault complexes with the property that it is compatible with the $\bar{\partial}$ -operator on both sides.

To see how this works, suppose $D : \mathcal{V}^{hol} \rightarrow \mathcal{W}^{hol}$ is locally of the form

$$D = \sum_{m_1, \dots, m_d} a_{m_1 \dots m_d}(z) \frac{\partial^{m_1}}{\partial z_1^{m_1}} \cdots \frac{\partial^{m_d}}{\partial z^{m_d}},$$

where $a_{m_1 \dots m_d}(z)$ denotes a local holomorphic section of $\text{Hom}(V, W)$. Then, if $\alpha = s_I(z, \bar{z}) d\bar{z}_I \in \Omega^{0,*}(X, V)$, where s_I is a local *smooth* section of V , we define

$$D^{\Omega^{0,*}} \alpha = \sum_{m_1, \dots, m_d} a_{m_1 \dots m_d}(z) \left(\frac{\partial^{m_1}}{\partial z_1^{m_1}} \cdots \frac{\partial^{m_d}}{\partial z^{m_d}} s_I(z, \bar{z}) \right) d\bar{z}_I \in \Omega^{0,*}(X, W).$$

In this way, D extends to a differential operator

$$D^{\Omega^{0,*}} : \Omega^{0,*}(X, V) \rightarrow \Omega^{0,*}(X, W).$$

Since D is holomorphic, it is immediate that $D^{\Omega^{0,*}}\bar{\partial}_V = \bar{\partial}_W D^{\Omega^{0,*}}$ where $\bar{\partial}_V, \bar{\partial}_W$ are the $(0,1)$ -connections on V, W respectively. Thus, $D^{\Omega^{0,*}}$ is a map of sheaves of cochain complexes.

Via this construction, we extend F_k to a $\Omega^{0,*}(X)$ -valued functional on $\Omega^{0,*}(X, V)$ by the formula

$$F_k^{\Omega^{0,*}} : (\alpha_1, \dots, \alpha_k) \mapsto \sum_{i_1, \dots, i_k} D_{i_1}^{\Omega^{0,*}}(\alpha_1) \wedge \dots \wedge D_{i_k}^{\Omega^{0,*}}(\alpha_k) \in \Omega^{0,*}(X).$$

Here, as above, the α_i 's denote sections in $\Omega^{0,*}(X, V)$.

We have thus produced a linear map

$$(-)^{\Omega^{0,*}} : \text{Hom}_{\mathcal{O}^{hol}}((J^{hol}V)^{\otimes k}, \mathcal{O}^{hol}) \rightarrow \text{Hom}_{C^\infty}((J\Omega^{0,*}(X, V))^{\otimes k}, \Omega^{0,*}(X))$$

where $J\Omega^{0,*}(X, V)$ denotes the sheaf of smooth jets of the graded vector bundle underlying the Dolbeault complex. This map clearly restricts to the symmetric coinvariants on both sides. Taking direct products and tensoring with $\Omega_X^{d, hol}$ we have a map

$$\begin{aligned} \text{Lag}^{hol}(V) &\rightarrow \Omega^{d, hol} \otimes_{C^\infty} \prod_{k>0} \text{Hom}_{C^\infty}(J\Omega^{0,*}(X, V)^{\otimes k}, \Omega^{0,*}(X)) \\ &\cong \Omega^{d,*} \otimes_{C^\infty} \prod_{k>0} \text{Hom}(J\Omega^{0,*}(X, V)^{\otimes k}, C^\infty). \end{aligned}$$

We have already mentioned that this map is compatible with the $\bar{\partial}$ -operator on the right-hand side. Moreover, the holomorphic differential operator Q^{hol} also extends to a differential operator on the right-hand side in a way compatible with $\bar{\partial}$. Thus, $(-)^{\Omega^{0,*}}$ is a map of cochain complexes, where $\text{Lag}^{hol}(X)$ is equipped with the differential Q^{hol} and the right-hand side has differential $\bar{\partial} + Q^{hol}$.

The right-hand side admits a map of degree $-d$ to $\Omega^{d,d} \otimes_{C^\infty} \prod_{k>0} \text{Hom}(J\Omega^{0,*}(X, V)^{\otimes k}, C^\infty)$ by projecting onto the (d, d) -component of $\Omega^{d,*}$. Note that this map is only graded linear, it does not preserve the $\bar{\partial}$ -differential. However, once we quotient by the action of vector fields we do get a well-defined map

$$\mathcal{O}_{loc}^{hol}(V) \rightarrow \Omega^{d,d} \otimes_D \prod_{k>0} \text{Hom}(J\Omega^{0,*}(X, V)^{\otimes k}, C^\infty)_{S_k}[-d].$$

Note that we have accounted for the shift of d coming from $\Omega^{d,*} \rightarrow \Omega^{d,d}[-d]$. The right-hand side is precisely the (shifted) space of ordinary local functionals for the sheaf $\mathcal{E}_V = \Omega^{0,*}(X, V)$ defined in Definition 2.1.

In conclusion, we have obtained the following map of sheaves of cochain complexes

$$(8) \quad \int (-)^{\Omega^{0,*}} : \mathcal{O}_{loc}^{hol}(V) \rightarrow \mathcal{O}_{loc}(\Omega^{0,*}(X, V))[-d].$$

In fact, we have the following stronger result, that this map is compatible with the brackets on both sides.

Lemma 2.20. *The map $\int(-)^{\Omega^{0,*}}$ defines an map of sheaves of dg Lie algebras*

$$\int(-)^{\Omega^{0,*}} : \mathcal{O}_{\text{loc}}^{\text{hol}}(V)[d-1] \rightarrow \mathcal{O}_{\text{loc}}(\mathcal{E}_V)[-1]$$

Proof. By definition, the sheaf of local functionals on \mathcal{E}_V is equal to

$$\text{Dens}_X \otimes_{D_X} \mathcal{O}_{\text{red}}(J\mathcal{E}_V).$$

Since $\mathcal{O}_{\text{red}}(J\mathcal{E}_V)$ is flat as a D_X -module [Cos11], we can replace the tensor product \otimes_{D_X} with the derived tensor product $\otimes_{D_X}^{\mathbb{L}}$.

We now use the following observation about D -modules. If M is a holomorphic D_X^{hol} -module, then clearly it forgets down to an ordinary smooth D_X -module (with the same underlying C_X^∞ -module structure) that we denote M^{C^∞} . Moreover, there is a quasi-isomorphism of D -modules

$$\Omega_X^{d,\text{hol}} \otimes_{D_X^{\text{hol}}}^{\mathbb{L}} M[d] \simeq \Omega_X^{d,d} \otimes_{D_X}^{\mathbb{L}} M^{\text{C}^\infty}.$$

We apply this to the case $M = \mathcal{O}_{\text{red}}(J^{\text{hol}}V)$, where V is a holomorphic vector bundle. This says that there is a quasi-isomorphism

$$(9) \quad \Omega_X^{d,\text{hol}} \otimes_{D_X^{\text{hol}}}^{\mathbb{L}} \mathcal{O}_{\text{red}}(J^{\text{hol}}V)[d] \simeq \Omega_X^{d,d} \otimes_{D_X}^{\mathbb{L}} \mathcal{O}_{\text{red}}(J^{\text{hol}}V).$$

This quasi-isomorphism is compatible with the Q^{hol} differential and the bracket $\{-, -\}^{\text{hol}}$ on both sides. Note that the left-hand side is simply the space of shifted holomorphic local functionals $\mathcal{O}_{\text{loc}}^{\text{hol}}(V)[d]$.

Next, observe that the map $(-)^{\Omega^{0,*}}$ determines a map of sheaves of cochain complexes

$$(10) \quad (-)^{\Omega^{0,*}} : \Omega_X^{d,d} \otimes_{D_X}^{\mathbb{L}} \mathcal{O}_{\text{red}}(J^{\text{hol}}V) \rightarrow \text{Dens}_X \otimes_{D_X}^{\mathbb{L}} \mathcal{O}_{\text{red}}(J\mathcal{E}_V).$$

The right-hand side is quasi-isomorphic to $\mathcal{O}_{\text{loc}}(\mathcal{E}_V)$. The composition of (9) and (10) is simply the map (8)

$$\int(-)^{\Omega^{0,*}} : \mathcal{O}_{\text{loc}}^{\text{hol}}(V)[d] \rightarrow \mathcal{O}_{\text{loc}}(\mathcal{E}_V).$$

One checks immediately that this map is compatible with the brackets, namely

$$\{I^{\text{hol}}, J^{\text{hol}}\}^{\text{hol}} = \left\{ \int I^{\Omega^{0,*}}, \int J^{\Omega^{0,*}} \right\}.$$

□

As a result of the equivalence between solutions to the classical master equation and Maurer-Cartan elements in the dg Lie algebras of shifted local functionals, we have the following.

Proposition 2.21. *Every classical holomorphic theory $(V, Q^{\text{hol}}, (-, -)_V, I^{\text{hol}})$ determines the structure of a classical BV theory. The underlying free BV theory is given in Definition/Lemma 1 $(\mathcal{E}_V, Q, \omega_V)$ and the interaction is $I = \int I^{\Omega^{0,*}}$.*

Table 1 is a useful summary showing how we are producing a BV theory from a holomorphic theory.

Holomorphic theory	BV theory
Holomorphic bundle V	Space of fields $\mathcal{E}_V = \Omega^{0,*}(X, V)$
Holomorphic differential operator Q^{hol}	Linear BRST operator $\bar{\partial} + Q^{hol}$
Non-degenerate pairing $(-, -)_V$	(-1) -symplectic structure ω_V
Holomorphic Lagrangian I^{hol}	Local functional $I = \int I^{\Omega^{0,*}} \in \mathcal{O}_{loc}(\mathcal{E}_V)$

TABLE 1. From holomorphic to BV

Example 2.22. Holomorphic BF-theory Let \mathfrak{g} be a Lie algebra and X any complex manifold. Consider the following holomorphic vector bundle on X :

$$V = \underline{\mathfrak{g}}_X[1] \oplus K_X \otimes \mathfrak{g}^\vee[d-2].$$

The notation $\underline{\mathfrak{g}}_X$ denotes the trivial bundle with fiber \mathfrak{g} . The pairing $V \otimes V \rightarrow K_X[d-1]$ is similar to the pairing for the $\beta\gamma$ system, except we use the evaluation pairing $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$ between \mathfrak{g} and its dual \mathfrak{g}^\vee . In this example, $Q^{hol} = 0$.

We describe the holomorphic Lagrangian. If $f_i : X \rightarrow \mathbb{C}, i = 1, 2$ are holomorphic functions and $\beta \in K_X$, consider the trilinear functional

$$I^{hol}(f_1 \otimes X_1, f_2 \otimes X_2, \beta \otimes X^\vee) = f_1 f_2 \beta \langle X^\vee, [X_1, X_2] \rangle_{\mathfrak{g}} + \dots$$

where the \dots means that we symmetrize the inputs. This defines an element $I^{hol} \in \mathcal{O}_{loc}^{hol}(V)^+$ and the Jacobi identity for \mathfrak{g} guarantees $\{I^{hol}, I^{hol}\}^{hol} = 0$. The fields of the corresponding BV theory are

$$\mathcal{E}_V = \Omega^{0,*}(X, \mathfrak{g})[1] \oplus \Omega^{d,*}(X, \mathfrak{g}^*)[d-2].$$

The induced local functional $I^{\Omega^{0,*}}$ on \mathcal{E}_V is

$$I^{\Omega^{0,*}}(\alpha, \beta) = \int_X \langle \beta, [\alpha, \alpha] \rangle_{\mathfrak{g}}.$$

The total action is $S(\alpha, \beta) = \int \langle \beta, \bar{\partial}\alpha \rangle + \langle \beta, [\alpha, \alpha] \rangle_{\mathfrak{g}}$. This is formally similar to *BF* theory (see below) and for that reason we refer to it as *holomorphic BF* theory. The moduli problem this describes is the cotangent theory to the moduli space of holomorphic connections on the trivial G -bundle near the trivial bundle. There is an obvious enhancement that works near any holomorphic principal bundle. When $d = 2$, in [Joh94], or for a more mathematical treatment see [Cosc], it is shown that this theory is a twist of $\mathcal{N} = 1$ supersymmetric pure Yang-Mills on \mathbb{R}^4 .

Example 2.23. Topological BF-theory This is a deformation of the previous example that has appeared throughout the physics literature. Suppose we take as our graded holomorphic vector bundle

$$V = \left(\underline{\mathfrak{g}}_X \otimes \left(\bigoplus_{k=0}^d \wedge^k T^{*1,0} X[1-k] \right) \right) \oplus \left(\underline{\mathfrak{g}}_X^* \otimes \left(\bigoplus_{k=0}^d \wedge^k T^{*1,0} X[2(d-1)-k] \right) \right).$$

Here $\wedge^0 T^{*1,0} X$ is understood as the trivial bundle $\underline{\mathbb{C}}_X$. The pairing is given by combining the evaluation pairing between \mathfrak{g} and \mathfrak{g}^* and taking the wedge product and projecting onto the components isomorphic to K_X . Explicitly, the pairing is equal to the sum of bundle maps of the form

$$\text{ev}_{\mathfrak{g}} \otimes \wedge : \left(\underline{\mathfrak{g}}_X \otimes \wedge^k T^{*1,0} X[1-k] \right) \otimes \left(\underline{\mathfrak{g}}^*_X \otimes \wedge^{d-k} T^{*1,0} X[d-1+k] \right) \rightarrow K_X[d-1].$$

The holomorphic differential is of the form

$$Q^{hol} = \text{id}_{\mathfrak{g}} \otimes \partial + \text{id}_{\mathfrak{g}^*} \otimes \partial,$$

where ∂ is the holomorphic de Rham differential. The holomorphic interaction is given by combining the Lie algebra structure on \mathfrak{g} with the wedge product of the holomorphic bundles $\wedge^k T^{*1,0} X$. We observe that the associated BV theory has classical space of fields given by

$$(A, B) \in \mathcal{E}_V = \Omega^*(X, \mathfrak{g}[1] \oplus \mathfrak{g}^*[2d-2])$$

where Ω^* is now the *full* de Rham complex. The action functional is

$$S = \int_X \langle B, dA \rangle_{\mathfrak{g}} + \frac{1}{3} \langle B, [A, A] \rangle_{\mathfrak{g}}.$$

As above, $\langle -, - \rangle_{\mathfrak{g}}$ denotes the pairing between \mathfrak{g} and its dual. This is the well-known topological BF theory on the even dimensional *real* manifold X (of real dimension $2d$). It might seem silly that we have used the formalism of holomorphic field theory to describe a very simple topological theory. We will discuss advantages of this approach at the end of the next section. In particular, the theory of regularization for holomorphic theories we will employ has peculiar consequences for renormalizing certain classes of topological theories such as topological BF theory.

Remark 2.24. We have mentioned an alternative formulation of classical field theory in terms of sheaves of L_{∞} algebras. Just as in the ordinary case we can formulate the data of a classical holomorphic theory in terms of sheaves of L_{∞} algebras. We will not do that here, but hope the idea of how to do so is clear.

Remark 2.25. Our definition of a holomorphic theory is compatible with the definition of a two-dimensional chiral conformal field theory given in [Li] when the complex dimension is $d = 1$.

2.3. Holomorphically translation invariant theories. When working on affine space one can ask for a theory to be invariant with respect to translations. In this section, we take a break from holomorphic theories defined on general complex manifolds to consider the affine manifold $\mathbb{C}^d = \mathbb{R}^{2d}$. We recall what a *holomorphically translation invariant* theory is, and state a general result about deformations for such theories. This particular class of theories has been discussed in Chapter 10 of [CG], and it is a special case of a general holomorphic theory as defined above.

Let V be a holomorphic vector bundle on \mathbb{C}^n and suppose we fix an identification of bundles

$$V \cong \mathbb{C}^d \times V_0$$

where V_0 is the fiber of V at $0 \in \mathbb{C}^d$. We want to consider a classical theory with space of fields given by $\Omega^{0,*}(\mathbb{C}^d, V) \cong \Omega^{0,*}(\mathbb{C}^d) \otimes_{\mathbb{C}} V_0$. Moreover, we want this theory to be invariant with respect to the group of translations on \mathbb{C}^d . Per usual, it is best to work with the corresponding Lie algebra of translations. Using the complex structure, we choose a presentation for the complex Lie algebra of translations given by

$$\mathbb{C}^{2d} \cong \text{span}_{\mathbb{C}} \left\{ \frac{\partial}{\partial z_i}, \frac{\partial}{\partial \bar{z}_i} \right\}_{1 \leq i \leq d}.$$

To define a theory, we need to fix a non-degenerate pairing on V . Moreover, we want this to be translation invariant. So, suppose

$$(11) \quad (-, -)_V : V \otimes V \rightarrow K_{\mathbb{C}^d}[d-1]$$

is a skew-symmetric bundle map that is equivariant for the Lie algebra of translations. The shift is so that the resulting pairing on the Dolbeault complex is of the appropriate degree. Here, equivariance means that for sections v, v' we have

$$\left(\frac{\partial}{\partial z_i} v, v' \right)_V = L_{\partial z_i} (v, v')_V$$

where the right-hand side denotes the Lie derivative applied to $(v, v')_V \in \Omega_{\mathbb{C}^d}^{d, hol}$. There is a similar relation for the anti-holomorphic derivatives. We obtain a \mathbb{C} -valued pairing on $\Omega_{\mathbb{C}^d}^{0,*}(\mathbb{C}^d, V)$ via integration:

$$(12) \quad \int_{\mathbb{C}^d} \circ(-, -)_V : \Omega_{\mathbb{C}^d}^{0,*}(\mathbb{C}^d, V) \otimes \Omega_{\mathbb{C}^d}^{0,*}(\mathbb{C}^d, V) \xrightarrow{\wedge \circ(-, -)_V} \Omega^{d,*}(\mathbb{C}^d) \xrightarrow{\int} \mathbb{C}.$$

The first arrow is the wedge product of forms combined with the pairing on V . The second arrow is only nonzero on forms of type $\Omega^{d,d}$. Clearly, integration is translation invariant, so that the composition is as well.

The pairing (12) together with the differential $\bar{\partial}$ are enough to define a free theory. However, it is convenient to consider a slightly generalized version of this situation. We want to allow deformations of the differential $\bar{\partial}$ on Dolbeault forms of the form

$$Q = \bar{\partial} + Q^{hol}$$

where Q^{hol} is a holomorphic differential operator of the form

$$(13) \quad Q^{hol} = \sum_I \frac{\partial}{\partial z^I} \mu_I$$

where I is some multi-index and $\mu_I : V \rightarrow V$ is a linear map of cohomological degree $+1$. Note that we have automatically written Q^{hol} in a way that it is translation invariant. Of course, for this differential to define a free theory there needs to be some compatibility with the pairing on V .

We can summarize this in the following definition, which should be viewed as a slight modification of a free theory to this translation invariant holomorphic setting.

Definition 2.26. A holomorphically translation invariant free BV theory is the data of a holomorphic vector bundle V together with

- (1) an identification $V \cong \mathbb{C}^d \times V_0$;
- (2) a translation invariant skew-symmetric pairing $(-, -)_V$ as in (11);
- (3) a holomorphic differential operator Q^{hol} as in (13);

such that the following conditions hold

- (1) the induced \mathbb{C} -valued pairing $\int \circ (-, -)_V$ is non-degenerate;
- (2) the operator Q^{hol} satisfies $(\bar{\partial} + Q^{hol})^2 = 0$ and is skew self-adjoint for the pairing:

$$\int (Q^{hol}v, v')_V = \pm \int (v, Q^{hol}v').$$

The first condition is required so that we obtain an actual (-1) -shifted symplectic structure on $\Omega^{0,*}(\mathbb{C}^d, V)$. The second condition implies that the derivation $Q = \bar{\partial} + Q^{hol}$ defines a cochain complex

$$\mathcal{E}_V = \left(\Omega^{0,*}(\mathbb{C}^d, V), \bar{\partial} + Q^{hol} \right),$$

and that Q is skew self-adjoint for the symplectic structure. Thus, in particular, \mathcal{E}_V together with the pairing define a free BV theory in the ordinary sense. In the usual way, we obtain the action functional via

$$S(\varphi) = \int (\varphi, (\bar{\partial} + Q^{hol})\varphi)_V.$$

Before going further, we will give a familiar example from the last section.

Example 2.27. The free $\beta\gamma$ system on \mathbb{C}^d . Consider the $\beta\gamma$ system with coefficients in any holomorphic vector bundle from Example 2.10 (and the remarks after it) specialized to the manifold $X = \mathbb{C}^d$. One immediately checks that this is a holomorphically translation invariant free theory.

2.3.1. *Translation invariant interactions.* Let's fix a general free holomorphically translation invariant theory $(V, (-, -)_V, Q^{hol})$ as above. We now define what a holomorphically translation invariant interacting theory is. Recall, translations span a $2d$ -dimensional abelian Lie algebra $\mathbb{C}^{2d} = \mathbb{C} \left\{ \frac{\partial}{\partial z_i}, \frac{\partial}{\partial \bar{z}_i} \right\}$. The first condition that an interaction be holomorphically translation invariant is that it be translation invariant, so invariant for this Lie algebra. The additional condition is a bit more involved.

Let $\bar{\eta}_i$ denote the operator on Dolbeault forms given by contraction with the antiholomorphic vector field $\frac{\partial}{\partial \bar{z}_i}$. Note that η_i acts on the Dolbeault complex on \mathbb{C}^d with values in any vector bundle. In particular it acts on the fields of a free holomorphically translation invariant theory as above, in addition to functionals on fields.

Definition 2.28. A holomorphically translation invariant local functional is a translation invariant local functional $I \in \mathcal{O}_{\text{loc}}(\mathcal{E}_V)^{\mathbb{C}^{2d}}$ such that $\bar{\eta}_i I = 0$ for all $1 \leq i \leq d$.

There is a succinct way of expressing holomorphic translation invariance as the Lie algebra invariants of a certain *dg Lie algebra*. Denote by $\mathbb{C}^d[1]$ the abelian d -dimensional graded Lie algebra concentrated in degree -1 by the elements $\{\bar{\eta}_i\}$. We want to consider deformations that are invariant for the action by the total *dg Lie algebra* $\mathbb{C}^{2d|d} = \mathbb{C}^{2d} \oplus \mathbb{C}^d[1]$. The differential sends $\eta_i \mapsto \frac{\partial}{\partial \bar{z}_i}$. The space of holomorphically translation invariant local functionals are denoted by $\mathcal{O}_{\text{loc}}(\mathcal{E}_V)^{\mathbb{C}^{2d|d}}$. The enveloping algebra of $\mathbb{C}^{2d|d}$ is of the form

$$U(\mathbb{C}^{2d|d}) = \mathbb{C} \left[\frac{\partial}{\partial z_i}, \frac{\partial}{\partial \bar{z}_i}, \eta_i \right]$$

with differential induced from that in $\mathbb{C}^{2d|d}$. Note that this algebra is quasi-isomorphic to the algebra of constant coefficient polynomial holomorphic differential operators $\mathbb{C}[\partial/\partial z_i] \xrightarrow{\cong} U(\mathbb{C}^{2d|d})$.

From the definitions, we see that any translation invariant local functional is a sum of functionals of the form

$$\varphi \mapsto \int_{\mathbb{C}^d} F(D_1 \alpha, \dots, D_k \alpha) d^d z$$

where D_α is an operator in the space

$$\mathbb{C} \left[d\bar{z}_i, \frac{\partial}{\partial z_i}, \frac{\partial}{\partial \bar{z}_i}, \eta_i \right],$$

and $F : \Omega^{0,*}(\mathbb{C}^d, V)^{\otimes k} \rightarrow \Omega^{0,*}(\mathbb{C}^d)$ is a linear map of the form

$$\Omega^{0,*}(\mathbb{C}^d, V)^{\otimes k} \cong (\Omega^{0,*}(\mathbb{C}^d) \otimes V_0)^{\otimes k} = \Omega^{0,*}(\mathbb{C}^d)^{\otimes k} \otimes V_0^{\otimes k} \xrightarrow{\wedge^{\otimes F_0}} \Omega^{0,*}(\mathbb{C}^d),$$

where $F_0 : V_0^{\otimes k} \rightarrow \mathbb{C}$ is a linear map and \wedge denotes the wedge product of forms.

The condition $\eta_i I = 0$ means that none of the D_i 's have any $d\bar{z}_j$ -dependence. Using this description we can exhibit the space of holomorphically translation functionals as follows. Note that if E is any vector bundle on \mathbb{C}^d we can consider the fiber at zero of its jet bundle that we denote $J_0 E$.

Lemma 2.29. Let V be a holomorphic vector bundle on \mathbb{C}^d and denote $\mathcal{E}_V = \Omega^{0,*}(X, V)$. Then

$$\mathcal{O}_{\text{loc}}(\mathcal{E}_V)^{\mathbb{C}^{2d|d}} \cong \mathbb{C} \cdot d^d z \otimes_{U(\mathbb{C}^{2d|d})} \mathcal{O}_{\text{red}}(J_0 E_V)$$

where E_V is the vector bundle on \mathbb{C}^d such that $\mathcal{E}_V = \Gamma(E_V)$.

This description of holomorphically translation invariant local functionals allows us to give a convenient description of deformations of holomorphically translation invariant theories. Suppose $(V, Q^{\text{hol}}, (-, -)_V, I)$ be the data of an interacting holomorphically translation invariant theory on \mathbb{C}^d . We have already encountered the space of local functionals $\mathcal{O}_{\text{loc}}(\mathcal{E}_V)$ and the deformation complex of the interacting BV theory is

$$\text{Def}_{\mathcal{E}_V} = \left(\mathcal{O}_{\text{loc}}(\mathcal{E}_V), \bar{\partial} + Q^{\text{hol}} + \{I, -\} \right).$$

We'd like to characterize deformations that preserve holomorphically translation invariance.

Recall that in the holomorphic case there is the holomorphic jet bundle $J^{hol}V$. The fiber at zero of this jet bundle may be identified as $J_0^{hol}V = V_0[[z_1, \dots, z_d]]$ where the z_i 's denote the formal jet coordinate.

Corollary 2.30. *Suppose that $Q^{hol} = 0$. Then, there is a quasi-isomorphism*

$$(\text{Def}_{\mathcal{E}_V})^{\mathbb{C}^{2d|d}} \simeq \mathbb{C} \cdot \mathfrak{d}^d z \otimes_{\mathbb{C}[\partial_{z_1}, \dots, \partial_{z_d}]}^{\mathbb{L}} \mathcal{O}_{red}(V_0[[z_1, \dots, z_d]])[d].$$

Equipped with differential $\{I^{hol}, -\}$ where I^{hol} only depends on holomorphic differential operators. Here, $\partial_{z_i} = \frac{\partial}{\partial z_i}$ and $\mathbb{C} \cdot \mathfrak{d}^d z$ denotes the trivial right $\mathbb{C}[\partial_{z_i}]$ -module.

The local functional I defining the classical holomorphic theory endows $J^{hol}V[-1]$ the structure of a L_∞ algebra in $D_{\mathbb{C}^d}$ -modules. Repackaging the statement using Lie algebraic data we can rewrite the equivalence in the lemma as

$$(\text{Def}_{\mathcal{E}_V})^{\mathbb{C}^{2d|d}} \simeq \mathbb{C} \cdot \mathfrak{d}^d z \otimes_{\mathbb{C}[\partial_{z_1}, \dots, \partial_{z_d}]}^{\mathbb{L}} \mathbb{C}_{\text{Lie,red}}^*(V_0[[z]][-1])[d].$$

Proof. By Lemma 2.29 we have an expression for the holomorphically translation local functionals

$$(\text{Def}_{\mathcal{E}_V})^{\mathbb{C}^{2d|d}} = \left(\mathbb{C} \cdot \mathfrak{d}^d z \otimes_{U(\mathbb{C}^{2d|d})} \mathcal{O}_{red}(J_0 E_V)[d], \bar{\partial} + \{I, -\} \right).$$

Since $\mathcal{O}_{red}(J_0 E_V)$ is flat as a $U(\mathbb{C}^{2d|d})$ -module, it follows that we can replace the tensor product by the derived tensor product $\otimes^{\mathbb{L}}$ up to quasi-isomorphism so that

$$(\text{Def}_{\mathcal{E}_V})^{\mathbb{C}^{2d|d}} \simeq \left(\mathbb{C} \cdot \mathfrak{d}^d z \otimes_{U(\mathbb{C}^{2d|d})}^{\mathbb{L}} \mathcal{O}_{red}(J_0 E_V)[d], \bar{\partial} + \{I, -\} \right).$$

Consider the complex $\left(\mathcal{O}_{red}(J_0 E_V), \bar{\partial} + \{I, -\} \right)$. This complex is graded by symmetric degree, and the associated spectral sequence has first page the associated graded of $\mathcal{O}_{red}(J_0 E_V)$ equipped with the $\bar{\partial}$ differential. Moreover, at the E_1 -page, we have the quasi-isomorphism

$$\left(\mathcal{O}(J_0 E_V), \bar{\partial} \right) = \left(\mathcal{O}_{red}(V_0[[z_i, \bar{z}_i]][d\bar{z}_i]), \bar{\partial} \right) \simeq \mathcal{O}_{red}(V_0[[z_i]]).$$

Finally, we have already remarked that there is a quasi-isomorphism of algebras $U(\mathbb{C}^{2d|d}) \simeq U(\mathbb{C}^d)$ where the right-hand site is generated by the constant holomorphic vector fields. The proof of the claim follows. □

Example 2.31. The interacting $\beta\gamma$ system Suppose V is a complex vector space and consider the free $\beta\gamma$ system with values in V . The fields are $(\gamma, \beta) \in \Omega^{0,*}(\mathbb{C}^d) \otimes V \oplus \Omega^{d,*}(\mathbb{C}^d) \otimes$

$V^*[d-1]$. A holomorphically translation invariant differential operator on \mathbb{C}^d is a linear combination of operators of the form

$$(14) \quad D = \frac{\partial^{|I|}}{\partial z_I}$$

where $I = (i_1, \dots, i_d)$ is a multi-index, $\partial z_I = \partial z_{i_1} \cdots \partial z_{i_d}$ and $|I| = i_1 + \cdots + i_d$.

We get a family of holomorphically translation invariant interactions in the following way. For simplicity, suppose $V = \mathbb{C}$. Let ϵ be an arbitrary translation invariant Dolbeault form of type $(0, 1)$. An example of a holomorphically translation invariant interaction for the $\beta\gamma$ system with values in V is

$$I(\beta, \gamma) = \int_{\mathbb{C}^d} \epsilon \wedge \langle \beta, (D_1\gamma) \cdots (D_k\gamma) \rangle_V$$

where D_i are holomorphically translation invariant differential operators, like in Equation (14), on \mathbb{C}^d .

Example 2.32. Holomorphic superpotential. This is a different flavor of a holomorphically translation invariant interaction for the $\beta\gamma$ system, which is largely motivated by physics. Consider again, the $\beta\gamma$ system on \mathbb{C}^d with values in V . In addition, let $W \in \mathbb{C}[V]$ be a polynomial on the vector space V . Then, W extends in a natural way to a Dolbeault valued functional on $\Omega^{0,*}(\mathbb{C}^d) \otimes V$. One defines the local functional

$$I_W(\beta, \gamma) = \int_{\mathbb{C}^d} d^d z W(\gamma).$$

It is immediate to see that I_W is holomorphically translation invariant. On the other hand, it is not, in general, a degree zero functional. Hence, it does not define a classical theory in the usual sense. It does, however, define a slightly weaker classical theory that is only $\mathbb{Z}/2$ graded rather than the usual \mathbb{Z} grading we are accustomed to.

We don't develop the formal definition here, but I_W defines a holomorphically translation invariant $\mathbb{Z}/2$ -graded BV theory. When $d = 2$, the $\beta\gamma$ system arises as the minimal twist of the free $\mathcal{N} = 1$ chiral supermultiplet on \mathbb{R}^4 . In the presence of the interaction I_W , the theory is equal to the minimal twist of the $\mathcal{N} = 1$ chiral multiplet with *holomorphic superpotential* given by W .³

3. ONE-LOOP REGULARIZATION FOR THEORIES ON \mathbb{C}^d

In Wilsonian's approach to quantum field theory, constructing the path integral involves exhibiting a family of theories parametrized by some scale $L > 0$, that we take for illustration to be in units of length. The main idea is that the theory at scale L describes all interactions happening at length scales smaller than or equal to L . To obtain the full perturbative QFT, one takes the limit $L \rightarrow \infty$, where all quantum interactions are included.

³In super language, the superpotential term is usually written as $\int d^2\theta \int d^4x W(\Phi)$, where Φ is the chiral superfield.

In practice, one has a good handle on the theory between some finite scales $\epsilon < L$, and to obtain the theory at scale L one must make sense of the $\epsilon \rightarrow 0$ limit. Generally speaking, the naive limit is ill-defined; this is the part of the strategy for constructing a QFT where renormalization comes in.

In this section we consider the renormalization of holomorphic field theories on \mathbb{C}^d , for general $d \geq 1$. We start with a classical holomorphic theory on \mathbb{C}^d and study its one-loop homotopy renormalization group flow from some finite scale ϵ to scale L . This is where the theory is completely well-defined. Explicitly, this flow manifests as a sum over weights of graphs; that is, *Feynman diagrams*. In terms of diagrams, we consider the sum over graphs of genus at most one where at each vertex we place the holomorphic interaction defining the classical theory. The edges of the graphs are labeled by the propagator, which, for us, is an effective replacement for the Green's function of the $\bar{\partial}$ operator defining the kinetic piece of the holomorphic field theory.

To obtain a quantization of a classical theory one must make sense of the $\epsilon \rightarrow 0$ limit of this construction. In general, this involves introducing a family of *counterterms*. The presence of counterterms can be an often undesirable, but necessary part of constructing a quantum field theory. On one hand, logarithmic counterterms encode the β -function of an interacting field theory, which is a sensitive invariant and is important quantity to experimentally measure quantities in QFT. Roughly, this quantity measures how couplings run with renormalization group flow. Counterterms can also be extremely unwieldy. For instance, some theories of gravity require the introduction of infinitely many such counterterms [tHV74]. In this paper, we show how holomorphic theories on flat space are as well-behaved as possible when it comes to renormalization.

Our main result in this section is the following (which we state more carefully in Theorem 3.4 below):

Theorem 3.1. *For a holomorphic theory on \mathbb{C}^d , there exists a one-loop (pre)quantization where the naive $\epsilon \rightarrow 0$ limit exists and no counterterms are required.*

Remark 3.2. Already, in [Li] Li has proved a stronger version of Theorem 3.1 when the complex dimension is $d = 1$. His result holds to all orders in \hbar , and applies it to give an elegant interpretation of the quantum master equation for chiral conformal field theories on (flat) Riemann surfaces using vertex algebras. Although we do not make any statements in this thesis past one-loop quantizations, the higher loop behavior remains a rich and subtle problem that we hope to return to.

As a peculiar corollary of our main result, and our work in developing the one-loop β -function for QFT in the BV formalism [EWY18], we have the following.

Corollary 3.3. *The one-loop β -function of a holomorphic theory on \mathbb{C}^d is identically zero.*

This corollary has “no-go” style consequences for twists of supersymmetric field theories. As we have already mentioned, often times a supersymmetric field theory on \mathbb{R}^{2d} admits a holomorphic twist where half of the translations are left Q -exact. This result implies that the β -function is not protected under such holomorphic twists. For instance, $\mathcal{N} = 1$ supersymmetric Yang-Mills on \mathbb{R}^4 admits a holomorphic twist to holomorphic BF theory. While Yang-Mills has a non-trivial β -function, our results show that the β -function for holomorphic BF theory is zero.

The proof of the main result will be involve explicit evaluations and estimates of weights of Feynman diagrams. Before proceeding with the core analysis, we set up the problem using our notation and conventions used above.

Suppose $(V, Q^{hol}, (-, -)_V)$ prescribes the data of a free holomorphic theory on \mathbb{C}^d . This means that V is a holomorphic bundle on \mathbb{C}^d , $Q^{hol} : \mathcal{V}^{hol} \rightarrow \mathcal{V}^{hol}$ is a holomorphic differential operator, and $(-, -)_V$ is a (shifted) $K_{\mathbb{C}^d}$ -valued pairing on V . We assume, in addition, that Q^{hol} is translation invariant. Concretely, this means that

$$Q^{hol} \in \mathbb{C} \left[\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_d} \right].$$

The complex of fields, in the BV formalism, are given by the following deformed Dolbeault complex

$$\mathcal{E}_V = \left(\Omega^{0,*}(\mathbb{C}^d, V), \bar{\partial} + Q^{hol} \right).$$

We will fix a trivialization for the holomorphic vector bundle $V = \mathbb{C}^d \times V_0$, where V_0 is the fiber over $0 \in \mathbb{C}^d$. This leads to an identification $\Omega^{0,*}(\mathbb{C}^d, V) = \Omega^{0,*}(\mathbb{C}^d) \otimes_{\mathbb{C}} V_0$. Further, we write the (-1) -shifted symplectic structure defining the classical BV theory in the form

$$\omega_V(\alpha \otimes v, \beta \otimes w) = (v, w)_{V_0} \int d^d z (\alpha \wedge \beta)$$

where $(-, -)_{V_0}$ is a degree $(d-1)$ -shifted pairing on the finite dimensional vector space V_0 .

A holomorphic interacting theory is prescribed by a holomorphic Lagrangian $I^{hol} \in \mathcal{O}_{\text{loc}}^{hol}(V)^+$. As we have seen in Section 2.2.2 any holomorphic Lagrangian determines a local functional on its Dolbeault complex via integration $I = \int_X I^{\Omega^{0,*}}$. Here, as above, the notation $I^{\Omega^{0,*}}$ denotes the canonical extension of I^{hol} to the Dolbeault complex for V . Using the trivialization $V = \mathbb{C}^d \times V_0$ and $\Omega^{d,hol} = \mathbb{C} \cdot d^d z$, we can express the local functional as

$$I_k(\alpha) = \int I_k^{hol}(\alpha) = \int D_{k,1}(\phi_{k,1}(\alpha)) \cdots D_{k,k}(\phi_{k,k}(\alpha)) d^d z$$

where each $D_{i,j}$ is a holomorphic differential operator $D_{i,j} \in \mathbb{C} \left[\frac{\partial}{\partial z_i} \right]$, and $\phi_{i,j} \in V_0^{\vee}$.

3.1. Homotopy RG flow. As we've already mentioned, the main goal of this section is to show that for holomorphic theories on \mathbb{C}^d the one-loop renormalization group flow produces a prequantization modulo \hbar^2 . We follow the terminology of [Cos11] and use *prequantization* to refer to an effective family of functionals satisfying renormalization group flow but not necessarily the quantum master equation. We will see consequences of our result for solving the quantum master equation modulo \hbar^2 in the next section.

The building block in Costello's approach to renormalization is an effective family of functionals $\{I[L]\}$ parametrized by a *length scale* $L > 0$. For each $L > 0$ the functional $I[L] \in \mathcal{O}(\mathcal{E})[[\hbar]]$ must satisfy various conditions, which are carefully stated in Definition 8.2.9.1 of [CG]. We will recall some key aspects that will be useful for our purposes. The main condition is a compatibility between the functionals $I[L]$ as one changes the length scale; this is referred to as *homotopy renormalization group (RG) flow*. The flow from scale $L > 0$ to $L' > 0$ is encoded by an invertible linear map

$$W(P_{L < L'}, -) : \mathcal{O}(\mathcal{E})[[\hbar]] \rightarrow \mathcal{O}(\mathcal{E})[[\hbar]]$$

defined as a sum over weights of graphs $W(P_{L < L'}, I) = \sum_{\Gamma} W_{\Gamma}(P_{L < L'}, I)$. Here, Γ denotes a graph, and the weight W_{Γ} is defined as follows. One labels the vertices of valence k by the k th homogenous component of the functional I . The edges of the graph are labeled by the propagator $P_{L < L'}$. The total weight is given by iterative contractions of the homogenous components of the interaction with the propagator. For a more precise definition see Chapter 2 of [Cos11].

The family of functionals $\{I[L]\}$ defining a quantization must satisfy the *RG flow equation*

$$I[L'] = W(P_{L < L'}, I[L])$$

for all $L < L'$. Given a classical interaction $I \in \mathcal{O}_{\text{loc}}(\mathcal{E})$, there is a natural way to attempt construct an effective family of functionals satisfying the RG flow equations. Indeed, it follows from elementary properties of the homotopy RG flow operator $W(P_{L < L'}, -)$ that if the functional

$$I[L] \text{ " = " } W(P_{0 < L}, I)$$

were to be well-defined for each $L > 0$, then the RG flow equations would automatically be satisfied for the collection $\{I[L]\}$. The problem is that this naive guess is ill-defined due to the distributional nature of the propagator $P_{0 < L}$. The approach of Costello is to introduce a small parameter $\epsilon > 0$ and to consider the limit of the functionals $W(P_{\epsilon < L}, I)$ as $\epsilon \rightarrow 0$. For most theories, this $\epsilon \rightarrow 0$ limit is ill-defined, but there always exist ϵ -dependent *counterterms* $I^{CT}(\epsilon)$ rendering the existence of the $\epsilon \rightarrow 0$ limit of $W(P_{\epsilon < L}, I - I^{CT}(\epsilon))$.

Our main goal in this section amounts to showing that the naive $\epsilon \rightarrow 0$ limit exists without the necessity to introduce counterterms. This is a salient feature of holomorphic theories on \mathbb{C}^d that we will take advantage of to characterize anomalies, for instance.

We will only consider quantizations defined modulo \hbar^2 . In this case, the homotopy RG flow takes the explicit form:

$$W(P_{\epsilon < L}^V, I) = \sum_{\Gamma} \frac{\hbar^{s(\Gamma)}}{|\text{Aut}(\Gamma)|} W_{\Gamma}(P_{\epsilon < L}^V, I).$$

The sum is over graphs of genus ≤ 1 and W_{Γ} is the weight associated to the graph Γ .

We can now state the main result of this section.

Theorem 3.4. *Let \mathcal{E} be a holomorphic theory on \mathbb{C}^d with classical interaction I^{cl} . Then, there exists a one-loop prequantization $\{I[L] \mid L > 0\}$ of I^{cl} involving no counterterms. That is, the $\epsilon \rightarrow 0$ limit of*

$$W(P_{\epsilon < L}, I) \pmod{\hbar^2} \in \mathcal{O}(\mathcal{E})[[\hbar]]/\hbar^2$$

exists. Moreover, if I is holomorphically translation invariant we can pick the family $\{I[L]\}$ to be holomorphically translation invariant as well.

3.2. Holomorphic gauge fixing. The next component of a prequantization is the choice of a gauge fixing condition. From a physics point of view the choice of a gauge fixing condition is common place when computing quantities in QFT. Mathematically, it is equivalent to choosing an isotropic subspace of the space of fields which is necessary to define the path integral in the BV formalism. In our philosophy of QFT, all theories are really defined over the space (or simplicial set) of gauge fixing conditions. The theory does not depend on a gauge fixing condition in the sense that a path in the space of gauge fixing conditions leads to a homotopy between the associated theories. See Chapter 5 of [Cos11] for a thorough formulation of this.

In our approach, a gauge fixing condition appears through the choice gauge fixing operator is a square-zero operator on fields

$$Q^{GF} : \mathcal{E}_V \rightarrow \mathcal{E}_V[-1],$$

of cohomological degree -1 such that $[Q, Q^{GF}]$ is a generalized Laplacian on \mathcal{E} where Q is the linearized BRST operator. For a complete definition see Section 8.2.1 of [CG].

For holomorphic theories there is a convenient choice for a gauge fixing operator. To construct it we fix the standard flat metric on \mathbb{C}^d . Doing this, we let $\bar{\partial}^*$ be the adjoint of the operator $\bar{\partial}$. Using the coordinates on $(z_1, \dots, z_d) \in \mathbb{C}^d$ we can write this operator as

$$\bar{\partial}^* = \sum_{i=1}^d \frac{\partial}{\partial(d\bar{z}_i)} \frac{\partial}{\partial z_i}.$$

The operator $\frac{\partial}{\partial(d\bar{z}_i)}$ is the contraction with the anti-holomorphic vector field $\frac{\partial}{\partial \bar{z}_i}$. The operator $\bar{\partial}^*$ extends to the complex of fields via the formula

$$Q^{GF} = \bar{\partial}^* \otimes \text{id}_V : \Omega^{0,*}(X, V) \rightarrow \Omega^{0,*-1}(X, V),$$

Lemma 3.5. *The operator $Q^{GF} = \bar{\partial}^* \otimes \text{id}_V$ is a gauge fixing operator for the free theory $(\mathcal{E}_V, \bar{\partial} + Q, \omega_V)$.*

Proof. Clearly, Q^{GF} is square zero since $(\bar{\partial}^*)^2 = 0$. Since Q^{hol} is a translation invariant holomorphic differential operator we have

$$[\bar{\partial} + Q^{hol}, Q^{GF}] = [\bar{\partial}, \bar{\partial}^*] \otimes \text{id}_V.$$

The operator $[\bar{\partial}, \bar{\partial}^*]$ is the Dolbeault Laplacian $\Delta_{\bar{\partial}}$ on \mathbb{C}^d , which in coordinates is

$$\Delta_{\bar{\partial}} = - \sum_{i=1}^d \frac{\partial}{\partial \bar{z}_i} \frac{\partial}{\partial z_i}.$$

In particular, the operator $[\bar{\partial}, \bar{\partial}^*] \otimes \text{id}_V$ is a generalized Laplacian.

Finally, we must show that Q^{GF} is (graded) self-adjoint for the shifted symplectic pairing ω_V . This follows from the fact about Dolbeault forms on \mathbb{C}^d . If $\alpha, \beta \in \Omega_c^{0,*}(\mathbb{C}^d)$ then

$$\int_{\mathbb{C}^d} (\bar{\partial}^* \alpha) \wedge \beta \wedge d^d z = \pm \int_{\mathbb{C}^d} \alpha \wedge (\bar{\partial}^* \beta) \wedge d^d z.$$

□

Remark 3.6. One may ask what happens if we choose a different metric on \mathbb{C}^d to define the gauge fixing operator. For every choice of a Hermitian metric h on \mathbb{C}^d we obtain an operator $\bar{\partial}_h^*$ and hence a gauge fixing condition. In fact, this defines a *family* of theories defined over the space of all Hermitian metrics. Since this space is affine, hence connected, we can always choose a path to the standard metric to any other one, thus resulting in a homotopy equivalence between prequantizations defined by the standard metric and the fixed one.

3.3. The propagator on \mathbb{C}^d . The gauge fixing operator determines a generalized Laplacian, which for us is essentially the ordinary Dolbeault Laplacian on \mathbb{C}^d . Our regularization scheme utilizes the heat kernel associated to the Laplacian, for which we recall the explicit form below. By definition, the scale $L > 0$ heat kernel is a symmetric element $K_L^V \in \mathcal{E}_V(\mathbb{C}^d) \otimes \mathcal{E}_V(\mathbb{C}^d)$ that satisfies

$$\omega_V(K_L, \varphi) = e^{-L[Q, Q^{GF}]} \varphi$$

for any field $\varphi \in \mathcal{E}_V$. Thus, it is an integral kernel for the operator $e^{-L[Q, Q^{GF}]}$. For a more detailed definition of how heat kernels are used to define a quantum field theory in the BV formalism, see Section 8.2.3 in [CG]. In this section we deduce the explicit form of the heat kernel for our holomorphic theory on \mathbb{C}^d .

The tensor square of $\mathcal{E}_V(\mathbb{C}^d)$ decomposes as

$$(15) \quad \mathcal{E}_V(\mathbb{C}^d) \otimes \mathcal{E}_V(\mathbb{C}^d) = \left(\Omega^{0,*}(\mathbb{C}^d) \otimes \Omega^{0,*}(\mathbb{C}^d) \right) \otimes (V_0 \otimes V_0).$$

We will decompose the heat kernel accordingly.

Pick a basis $\{e_i\}$ of V_0 and let

$$\mathbf{C}_{V_0} = \sum_{i,j} \omega_{ij}(e_i \otimes e_j) \in V_0 \otimes V_0$$

be the quadratic Casimir. Here, (ω_{ij}) is the inverse matrix to the pairing $(-, -)_{V_0}$.

Due to the nature of our symplectic pairing, we see that the heat kernel splits with respect to the decomposition in Equation (15) as

$$K_L^V(z, w) = K_L^{an}(z, w) \cdot \mathbf{C}_{V_0}.$$

The analytic part K_L^{an} is independent of V and equal to the heat kernel for Dolbeault Laplacian $\Delta_{\bar{\partial}}$ acting on Dolbeault forms on \mathbb{C}^d .

We can further split this analytic heat kernel as the heat kernel for the ordinary Laplacian acting on functions. Indeed, for $L > 0$ the analytic heat kernel K_L^{an} is equal to

$$K_L^{an}(z, w) = k_L^{an}(z, w) \prod_{i=1}^d (d\bar{z}_i - d\bar{z}_j) \in \Omega^{0,*}(\mathbb{C}^d) \otimes \Omega^{0,*}(\mathbb{C}^d) \cong \Omega^{0,*}(\mathbb{C}^d \times \mathbb{C}^d) \cong C^\infty(\mathbb{C}^d \times \mathbb{C}^d)[dz, dw]$$

where $k_L^{an}(z, w) \in C^\infty(\mathbb{C}^d \times \mathbb{C}^d)$ is the heat kernel for the Laplacian acting on functions. It is normalized by the rule

$$(e^{-L\Delta_{\bar{\partial}}} f)(z) = \int_{w \in \mathbb{C}^d} d^{2d}w k_L^{an}(z, w) f(w)$$

where $f \in C^\infty(\mathbb{C}^d)$. Explicitly, k_L^{an} is given by

$$k_L^{an}(z, w) = \frac{1}{(2\pi i L)^d} e^{-|z-w|^2/4L}.$$

The propagator for the holomorphic theory \mathcal{E}_V is defined using the heat kernels above by the equation

$$P_{\epsilon < L}^V(z, w) = \int_{t=\epsilon}^L dt (Q^{GF} \otimes 1) K_L^V(z, w).$$

Since the element \mathbf{C}_{V_0} is independent of the coordinate on \mathbb{C}^d , the propagator also decomposes as

$$P_{\epsilon < L}^V(z, w) = P_{\epsilon < L}^{an}(z, w) \cdot \mathbf{C}_{V_0}$$

where

$$\begin{aligned} P_{\epsilon < L}^{an}(z, w) &= \int_{t=\epsilon}^L dt (\bar{\partial}^* \otimes 1) K_L^V(z, w) \\ &= \int_{t=\epsilon}^L dt \frac{1}{(2\pi i t)^d} \sum_{j=1}^d (-1)^{j-1} \left(\frac{\bar{z}_j - \bar{w}_j}{4t} \right) e^{-|z-w|^2/4t} \prod_{i \neq j}^d (d\bar{z}_i - d\bar{z}_j). \end{aligned}$$

Remark 3.7. In the limit as $\epsilon \rightarrow 0$ and $L \rightarrow \infty$, this propagator reduces to the Green's function for the Dolbeault operator on \mathbb{C}^d . We can see this simplification explicitly.

First, we recall the form of the Green's function. Introduce the δ -distribution δ_Δ along the diagonal in $\mathbb{C}^d \times \mathbb{C}^d$. In formulas

$$\delta_\Delta : \Omega_c^{0,*}(\mathbb{C}^d) \times \Omega_c^{0,*} \rightarrow \mathbb{C} , \quad (\alpha, \beta) \mapsto \int_{\Delta \subset \mathbb{C}^d \times \mathbb{C}^d} \mathbf{d}^d z \wedge \alpha \wedge \beta.$$

The Green's function for $\bar{\partial}$ is given in terms of the *Bochner-Martinelli kernel*. To define it, first consider the smooth form on $\mathbb{C}^d \times \mathbb{C}^d$ away from the diagonal $\omega_{BM} \in \bar{\Omega}^{0,*}(\mathbb{C}_z^d \times \mathbb{C}_w^d \setminus \Delta)$ given by

$$\omega_{BM}(z, w) = \frac{(d-1)!}{(2\pi i)^d} \frac{1}{|z-w|^{2d}} \sum_{i=1}^d (-1)^{i-1} (\bar{z}_i - \bar{w}_i) \prod_{j \neq i} (\mathbf{d}\bar{z}_j - \mathbf{d}\bar{w}_j).$$

Since $\bar{\partial}\omega_{BM}(z, w) = 0$ away from the diagonal, we see that $\bar{\partial}\omega_{BM}$ extends to a distribution form on $\mathbb{C}^d \times \mathbb{C}^d$. Indeed, this distribution solves Green's equation for the $\bar{\partial}$ -operator

$$\bar{\partial}\omega_{BM} = \delta_\Delta.$$

For more details on the above kernel and its relation to higher residues, see Chapter 3 of [GH94] for instance. For now, we have the immediate calculation.

Lemma 3.8. *The $\epsilon \rightarrow 0, L \rightarrow \infty$ distributional limit of the propagator $P_{\epsilon < L}(z, w)$ exists. Moreover, as distributions*

$$\lim_{\epsilon \rightarrow 0} \lim_{L \rightarrow \infty} P_{\epsilon < L}(z, w) = \omega_{BM}(z, w).$$

Proof. Note that

$$\begin{aligned} P_{\epsilon < L}(z, w) &= \int_{t=\epsilon}^L \mathbf{d}t e^{-|z-w|^2/4t} \frac{1}{(2\pi i t)^d} \sum_{j=1}^d (-1)^{j-1} \frac{\bar{z}_j - \bar{w}_j}{4t} (\mathbf{d}^d z - \mathbf{d}^d w) \prod_{i \neq j} (\mathbf{d}\bar{z}_i - \mathbf{d}\bar{w}_i) \\ &= \frac{1}{(2\pi i)^d} \frac{1}{|z-w|^{2d}} \sum_j (-1)^{j-1} (\bar{z}_j - \bar{w}_j) (\mathbf{d}^d z - \mathbf{d}^d w) \\ &\quad \times \prod_{i \neq j} (\mathbf{d}\bar{z}_i - \mathbf{d}\bar{w}_i) \int_{u=|z-w|^2/L}^{|z-w|^2/\epsilon} \mathbf{d}u u^{d-1} e^{-u}. \end{aligned}$$

In the second line we have made the substitution $u = |z-w|^2/4t$. Integration over u produces the desired result. \square

Thus, the propagator $P_{\epsilon < L}$ is an effective replacement for the Green's function for $\bar{\partial}$ on \mathbb{C}^d .

3.4. Trees. We now turn to studying the one-loop effective action for the holomorphic theory on \mathbb{C}^d . For the genus zero graphs, or trees, we do not have any analytic difficulties to worry about. The propagator $P_{\epsilon < L}^V$ is smooth so long as $\epsilon, L > 0$ but when $\epsilon \rightarrow 0$ it inherits a singularity along the diagonal $z = w$. This is what contributes to the divergences in the naive definition of RG flow $W(P_{0 < L}, -)$. But, if Γ is a tree the weight $W_\Gamma(P_{0 < L}^V, I)$ only involves multiplication of distributions with transverse singular support, so is well-defined. Thus we have observed the following.

Lemma 3.9. *If Γ is a tree then $\lim_{\epsilon \rightarrow 0} W_\Gamma(P_{\epsilon < L}, I)$ exists.*

The only possible divergences in the $\epsilon \rightarrow 0$ limit, then, must come from graphs of genus one, which we now direct our attention to.

3.5. A simplification for one-loop weights. Every graph of genus one is a wheel with some trees protruding from the external edges of the tree. Thus, we can write the weight of a genus one graph as a product of weights associated to trees times the weight associated to a wheel. We have just observed that the weights associated to trees are automatically convergent in the $\epsilon \rightarrow 0$ limit, thus it suffices to focus on genus one graphs that are purely wheels with some number of external edges.

The definition of the weight of the wheel involves placing the propagator at each internal edge and the interaction I at each vertex. The weights are evaluated by placing compactly supported fields $\varphi \in \mathcal{E}_{V,c} = \Omega_c^{0,*}(\mathbb{C}^d, V)$ at each of the external edges. We will make two simplifications:

- (1) the only ϵ dependence appears in the analytic part of the propagator $P_{\epsilon < L}^{an}$, so we can forget about the combinatorial factor C_{V_0} and assume all external edges are labeled by compactly supported Dolbeault forms in $\Omega_c^{0,*}(\mathbb{C}^d)$;
- (2) each vertex labeled by I is a sum of interactions of the form

$$\int_{\mathbb{C}^d} D_1(\varphi) \cdots D_k(\varphi) d^d z$$

where D_i is a holomorphic differential operator (only involves $\frac{\partial}{\partial z_i}$ -derivatives). Some of the differential operators will hit the compactly supported Dolbeault forms placed on the external edges of the graph. The remaining operators will hit the internal edges labeled by the propagators. Since a holomorphic differential operator preserves the space of compactly supported Dolbeault forms that is independent of ϵ , we replace each input by an arbitrary compactly supported Dolbeault form.

Thus, for the $\epsilon \rightarrow 0$ behavior it suffices to look at weights of wheels with arbitrary compactly supported functions as inputs where each of the internal edges are labeled by some translation invariant holomorphic differential operator

$$D = \sum_{n_1, \dots, n_d} \frac{\partial^{n_1}}{\partial z_1^{n_1}} \cdots \frac{\partial^{n_d}}{\partial z_d^{n_d}}$$

applied to the propagator $P_{\epsilon < L}^{an}$. This motivates the following definition.

Definition 3.10. Let $\epsilon, L > 0$. In addition, fix the following data.

- (a) An integer $k \geq 1$ that will be the number of vertices of the graph.
- (b) For each $\alpha = 1, \dots, k$ a sequence of integers

$$\vec{n}^\alpha = (n_1^\alpha, \dots, n_d^\alpha).$$

We denote by $(\vec{n}) = (n_i^j)$ the corresponding $d \times k$ matrix of integers.

The analytic weight associated to the pair $(k, (\vec{n}))$ is the smooth distribution

$$W_{\epsilon < L}^{k, (\vec{n})} : C_c^\infty((\mathbb{C}^d)^k) \rightarrow \mathbb{C},$$

that sends a smooth compactly supported function $\Phi \in C_c^\infty((\mathbb{C}^d)^k) = C_c^\infty(\mathbb{C}^{dk})$ to

$$(16) \quad W_{\epsilon < L}^{k, (\vec{n})}(\Phi) = \int_{(z^1, \dots, z^k) \in (\mathbb{C}^d)^k} \prod_{\alpha=1}^k d^d z^\alpha \Phi(z^1, \dots, z^k) \prod_{\alpha=1}^k \left(\frac{\partial}{\partial z^\alpha} \right)^{\vec{n}^\alpha} P_{\epsilon < L}^{an}(z^\alpha, z^{\alpha+1}).$$

In the above expression, we use the convention that $z^{k+1} = z^1$.

The coordinate on $(\mathbb{C}^d)^k$ is given by $\{z_i^\alpha\}$ where $\alpha = 1, \dots, k$ and $i = 1, \dots, d$. For each α , $\{z_1^\alpha, \dots, z_d^\alpha\}$ is the coordinate for the space \mathbb{C}^d sitting at the vertex labeled by α . We have also used the shorthand notation

$$\left(\frac{\partial}{\partial z^\alpha} \right)^{\vec{n}^\alpha} = \frac{\partial^{n_1^\alpha}}{\partial z_1^\alpha} \cdots \frac{\partial^{n_d^\alpha}}{\partial z_d^\alpha}.$$

We will refer to the collection of data $(k, (\vec{n}))$ in the definition as *wheel data*. The motivation for this is that the weight $W_{\epsilon < L}^{k, (\vec{n})}$ is the analytic part of the full weight $W_\Gamma(P_{\epsilon < L}^V, I)$ where Γ is a wheel with k vertices.

We have reduced the proof of Proposition 3.4 to showing that the $\epsilon \rightarrow 0$ limit of the analytic weight $W_{\epsilon < L}^{k, (\vec{n})}(\Phi)$ exists for any choice of wheel data $(k, (\vec{n}))$. To do this, there are two steps. First, we show a vanishing result that says when $k \geq d$ the weights vanish for purely algebraic reasons. The second part is the most technical aspect of the chapter where we show that for $k > d$ the weights have nice asymptotic behavior as a function of ϵ .

Lemma 3.11. *Let $(k, (\vec{n}))$ be a pair of wheel data. If the number of vertices k satisfies $k \leq d$ then*

$$W_{\epsilon < L}^{k, (\vec{n})} = 0$$

as a distribution on \mathbb{C}^{dk} for any $\epsilon, L > 0$.

Proof. In the integral expression for the weight (16) there is the following factor involving the product over the edges of the propagators:

$$(17) \quad \prod_{\alpha=1}^k \left(\frac{\partial}{\partial z^\alpha} \right)^{\vec{n}^\alpha} P_{\epsilon < L}^{an}(z^\alpha, z^\alpha).$$

We will show that this expression is identically zero. To simplify the expression we first make the following change of coordinates on \mathbb{C}^{dk} :

$$(18) \quad w^\alpha = z^{\alpha+1} - z^\alpha, \quad 1 \leq \alpha < k$$

$$(19) \quad w^k = z^k.$$

Introduce the following operators

$$\eta^\alpha = \sum_{i=1}^d \bar{w}_i^\alpha \frac{\partial}{\partial(\bar{d}\bar{w}_i^\alpha)}$$

acting on differential forms on \mathbb{C}^{dk} . The operator η^α lowers the anti-holomorphic Dolbeault type by one : $\eta : (p, q) \rightarrow (p, q - 1)$. Equivalently, η^α is contraction with the anti-holomorphic Euler vector field $\bar{w}_i^\alpha \partial / \partial \bar{w}_i^\alpha$.

Once we do this, we see that the expression (17) can be written as

$$\left(\left(\sum_{\alpha=1}^{k-1} \eta^\alpha \right) \prod_{i=1}^d \left(\sum_{\alpha=1}^{k-1} \bar{d}\bar{w}_i^\alpha \right) \right) \prod_{\alpha=1}^{k-1} \left(\eta^\alpha \prod_{i=1}^d \bar{d}\bar{w}_i^\alpha \right).$$

Note that only the variables \bar{w}_i^α for $i = 1, \dots, d$ and $\alpha = 1, \dots, k - 1$ appear. Thus we can consider it as a form on $\mathbb{C}^{d(k-1)}$. As such a form it is of Dolbeault type $(0, (d - 1) + (k - 1)(d - 1)) = (0, (d - 1)k)$. If $k < d$ then clearly $(d - 1)k > d(k - 1)$ so the form has greater degree than the dimension of the manifold and hence it vanishes.

The case left to consider is when $k = d$. In this case, the expression in (17) can be written as

$$(20) \quad \left(\left(\sum_{\alpha=1}^{d-1} \eta^\alpha \right) \prod_{i=1}^d \left(\sum_{\alpha=1}^{d-1} \bar{d}\bar{w}_i^\alpha \right) \right) \prod_{\alpha=1}^{d-1} \left(\eta^\alpha \prod_{i=1}^d \bar{d}\bar{w}_i^\alpha \right).$$

Again, since only the variables \bar{w}_i^α for $i = 1, \dots, d$ and $\alpha = 1, \dots, d - 1$ appear, we can view this as a differential form on $\mathbb{C}^{d(d-1)}$. Furthermore, it is a form of type $(0, d(d - 1))$. For any vector field X on $\mathbb{C}^{d(d-1)}$ the interior derivative i_X is a graded derivation. Suppose ω_1, ω_2 are two $(0, *)$ forms on $\mathbb{C}^{d(d-1)}$ such that the sum of their degrees is equal to d^2 . Then, $\omega_1 \iota_X \omega_2$ is a top form for any vector field on $\mathbb{C}^{d(d-1)}$. Since $\omega_1 \omega_2 = 0$ for form type reasons, we conclude that $\omega_1 \iota_X \omega_2 = \pm (i_X \omega_1) \omega_2$ with sign depending on the dimension d . Applied to the vector field $\bar{z}_i^1 \partial / \partial \bar{w}_i^1$ in (20) we see that the expression can be written (up to a sign) as

$$\eta^1 \left(\sum_{\alpha=1}^{d-1} \eta^\alpha \prod_{i=1}^d \left(\sum_{\alpha=1}^{d-1} \bar{d}\bar{w}_i^\alpha \right) \right) \left(\prod_{i=1}^d \bar{d}\bar{w}_i^1 \right) \prod_{\alpha=2}^{d-1} \left(\eta^\alpha \prod_{i=1}^d \bar{d}\bar{w}_i^\alpha \right).$$

Repeating this, for $\alpha = 2, \dots, k - 1$ we can write this expression (up to a sign) as

$$\left(\eta_{k-1} \cdots \eta_2 \eta_1 \sum_{\alpha=1}^{k-1} \eta^\alpha \prod_{i=1}^d \left(\sum_{\alpha=1}^{k-1} \bar{d}\bar{w}_i^\alpha \right) \right) \prod_{\alpha=1}^{k-1} \prod_{i=1}^d \bar{d}\bar{w}_i^\alpha$$

The expression inside the parentheses is zero since each term in the sum over α involves a term like $\eta^\beta \eta^\beta = 0$. This completes the proof for $k = d$. \square

Lemma 3.12. *Let $(k, (\vec{n}))$ be a pair of wheel data such that $k > d$. Then the $\epsilon \rightarrow 0$ limit of the analytic weight*

$$\lim_{\epsilon \rightarrow 0} W_{\epsilon < L}^{k, (n)}$$

exists as a distribution on \mathbb{C}^{dk} .

Proof. We will bound the absolute value of the weight in Equation (16) and show that it has a well-defined $\epsilon \rightarrow 0$ limit. First, consider the change of coordinates as in Equations (18),(19). For any compactly supported function Φ we see that $W_{\epsilon < L}^{k,(n)}(\Phi)$ has the form (21)

$$\int_{w^k \in \mathbb{C}^d} d^d w^k \int_{(w_1, \dots, w_{k-1}) \in (\mathbb{C}^d)^{k-1}} \left(\prod_{\alpha=1}^{k-1} d^d w^\alpha \right) \Phi(w^1, \dots, w^k) \left(\prod_{\alpha=1}^{k-1} \left(\frac{\partial}{\partial w^\alpha} \right)^{\bar{n}^\alpha} P_{\epsilon < L}^{an}(w^\alpha) \right) \times \sum_{\alpha=1}^{k-1} \left(\frac{\partial}{\partial w^\alpha} \right)^{\bar{n}^k} P_{\epsilon < L}^{an} \left(\sum_{\alpha=1}^{k-1} w^\alpha \right).$$

For $\alpha = 1, \dots, k-1$ the notation $P_{\epsilon < L}^{an}(w^\alpha)$ makes sense since $P_{\epsilon < L}^{an}(z^\alpha, z^{\alpha+1})$ is only a function of $w^\alpha = z^{\alpha+1} - z^\alpha$. Similarly $P_{\epsilon < L}^{an}(z^{k+1}, z^1)$ is a function of

$$z^k - z^1 = \sum_{\alpha=1}^{k-1} w^\alpha.$$

Expanding out the propagators the weight takes the form

$$\int_{w^k \in \mathbb{C}^d} d^{2d} w^k \int_{(w_1, \dots, w_{k-1}) \in (\mathbb{C}^d)^{k-1}} \left(\prod_{\alpha=1}^{k-1} d^{2d} w^\alpha \right) \Phi(w^1, \dots, w^k) \int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} \prod_{\alpha=1}^k \frac{dt_\alpha}{(2\pi i t_\alpha)^d} \times \sum_{i_1, \dots, i_{k-1}=1}^d \epsilon_{i_1 \dots i_k} \left(\frac{\bar{w}_{i_1}^1 (\bar{w}^1)^{n^1}}{4t_1 |4t_1|^{n^1}} \right) \dots \left(\frac{\bar{w}_{i_{k-1}}^{k-1} (\bar{w}^{k-1})^{n^{k-1}}}{4t_{k-1} |4t_{k-1}|^{n^{k-1}}} \right) \left(\sum_{\alpha=1}^{k-1} \frac{\bar{w}_{i_k}^\alpha}{4t_k} \cdot \frac{1}{t^{|n^k|}} \left(\sum_{\alpha=1}^{k-1} \bar{w}^\alpha \right)^{n^k} \right) \times \exp \left(- \sum_{\alpha=1}^{k-1} \frac{|w^\alpha|^2}{4t_\alpha} - \frac{1}{4t_k} \left| \sum_{\alpha=1}^{k-1} w^\alpha \right|^2 \right)$$

The notation used above warrants some explanation. Recall, for each α the vector of integers is defined as $n^\alpha = (n_1^\alpha, \dots, n_d^\alpha)$. We use the notation

$$(\bar{w}^\alpha)^{n^\alpha} = \bar{w}_1^{n_1^\alpha} \dots \bar{w}_d^{n_d^\alpha}.$$

Furthermore, $|n^\alpha| = n_1^\alpha + \dots + n_d^\alpha$. Each factor of the form $\frac{\bar{w}_{i_\alpha}^\alpha}{t_\alpha}$ comes from the application of the operator $\frac{\partial}{\partial z_i}$ in $\bar{\partial}^*$ applied to the propagator. The factor $\frac{(\bar{w}^\alpha)^{n^\alpha}}{t^{|n^\alpha|}}$ comes from applying the operator $\left(\frac{\partial}{\partial w} \right)^{n^\alpha}$ to the propagator. Note that $\bar{\partial}^*$ commutes with any translation invariant holomorphic differential operator, so it doesn't matter which order we do this.

To bound this integral we will recognize each of the factors

$$\frac{\bar{w}_{i_\alpha}^\alpha (\bar{w}^\alpha)^{n^\alpha}}{4t_\alpha |4t_\alpha|^{n^\alpha}}$$

as coming from the application of a certain holomorphic differential operator to the exponential in the last line. We will then integrate by parts to obtain a simple Gaussian integral which will give us the necessary bounds in the t -variables. Let us denote this

Gaussian factor by

$$E(w, t) := \exp \left(- \sum_{\alpha=1}^{k-1} \frac{|w^\alpha|^2}{4t_\alpha} - \frac{1}{4t_k} \left| \sum_{\alpha=1}^{k-1} w^\alpha \right|^2 \right)$$

For each α, i_α introduce the $t = (t_1, \dots, t_k)$ -dependent holomorphic differential operator

$$D_{\alpha, i_\alpha}(t) := \left(\frac{\partial}{\partial w_{i_\alpha}^\alpha} - \sum_{\beta=1}^{k-1} \frac{t_\beta}{t_1 + \dots + t_k} \frac{\partial}{\partial w_{i_\alpha}^\beta} \right) \prod_{j=1}^d \left(\frac{\partial}{\partial w_j^\alpha} - \sum_{\beta=1}^{k-1} \frac{t_\beta}{t_1 + \dots + t_k} \frac{\partial}{\partial w_j^\beta} \right)^{n_j^\alpha}.$$

The following lemma is an immediate calculation

Lemma 3.13. *One has*

$$D_{\alpha, i_\alpha} E(w, t) = \frac{\bar{w}_{i_\alpha}^\alpha (\bar{w}^\alpha)^{n^\alpha}}{4t_\alpha t^{|\mathbf{n}^\alpha|}} E(w, t).$$

Note that all of the D_{α, i_α} operators mutually commute. Thus, we can integrate by parts iteratively to obtain the following expression for the weight:

$$\begin{aligned} & \pm \int_{w^k \in \mathbb{C}^d} d^{2d} w^k \int_{(w_1, \dots, w_{k-1}) \in (\mathbb{C}^d)^{k-1}} \left(\prod_{\alpha=1}^{k-1} d^{2d} w^\alpha \right) \int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} \prod_{\alpha=1}^k \frac{dt_\alpha}{(2\pi i t_\alpha)^d} \\ & \times \left(\sum_{i_1, \dots, i_k} \epsilon_{i_1 \dots i_d} D_{1, i_1} \cdots D_{k-1, i_{k-1}} \sum_{\alpha=1}^{k-1} D_{\alpha, i_\alpha} \Phi(w^1, \dots, w^k) \right) \\ & \times \exp \left(- \sum_{\alpha=1}^{k-1} \frac{|w^\alpha|^2}{t_\alpha} - \frac{1}{t_k} \left| \sum_{\alpha=1}^{k-1} w^\alpha \right|^2 \right). \end{aligned}$$

Thus, the absolute value of the weight is bounded by

$$(22) \quad \begin{aligned} |W_{\epsilon < L}^{k, (n)}(\Phi)| & \leq C \int_{w^k \in \mathbb{C}^d} d^{2d} w^k \int_{(w^1, \dots, w^{k-1}) \in (\mathbb{C}^d)^{k-1}} \prod_{\alpha=1}^{k-1} d^{2d} w^\alpha \Psi(w^1, \dots, w^{k-1}, w^k) \\ & \times \int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} dt_1 \dots dt_k \frac{1}{(4\pi)^{dk}} \frac{1}{t_1^d \dots t_k^d} \times E(w, t) \end{aligned}$$

where Ψ is some compactly supported function on \mathbb{C}^{dk} that is independent of t .

To compute the right hand side we will perform a Gaussian integration with respect to the variables (w^1, \dots, w^{k-1}) . To this end, notice that the exponential can be written as

$$E(w, t) = \exp \left(- \frac{1}{4} M_{\alpha\beta} (w^\alpha, w^\beta) \right)$$

where $(M_{\alpha\beta})$ is the $(k-1) \times (k-1)$ matrix given by

$$\begin{pmatrix} a_1 & b & b & \cdots & b \\ b & a_2 & b & \cdots & b \\ b & b & a_3 & \cdots & b \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ b & b & b & \cdots & a_{k-1} \end{pmatrix}$$

where $a_\alpha = t_\alpha^{-1} + t_k^{-1}$ and $b = t_k^{-1}$. The pairing (w^α, w^β) is the usual Hermitian pairing on \mathbb{C}^d , $(w^\alpha, w^\beta) = \sum_i w_i^\alpha \bar{w}_i^\beta$. After some straightforward linear algebra we find that

$$\det(M_{\alpha\beta})^{-1} = \frac{t_1 \cdots t_k}{t_1 + \cdots + t_k}.$$

We now perform a Wick expansion for the Gaussian integral in the variables (w^1, \dots, w^{k-1}) . For a reference similar to the notation used here see the Appendix of our work in [EWY18]. The inequality in (22) becomes

$$(23) \quad |W_{\epsilon < L}^{k,(n)}(\Phi)| \leq C' \int_{w^k \in \mathbb{C}^d} d^{2d} w^k \Psi(0, \dots, 0, w^k)$$

$$(24) \quad \times \int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} dt_1 \dots dt_k \frac{1}{(4\pi)^{dk}} \frac{1}{(t_1 \cdots t_k)^d} \left(\frac{t_1 \cdots t_k}{t_1 + \cdots + t_k} \right)^d + O(\epsilon)$$

$$(25) \quad = C' \int_{w^k \in \mathbb{C}^d} d^{2d} w^k \Psi(0, \dots, 0, w^k)$$

$$(26) \quad \times \int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} dt_1 \dots dt_k \frac{1}{(4\pi)^{dk}} \frac{1}{(t_1 + \cdots + t_k)^d} + O(\epsilon).$$

The first term in the Wick expansion is written out explicitly. The $O(\epsilon)$ refers to higher terms in the Wick expansion, which one can show all have order ϵ , so disappear in the $\epsilon \rightarrow 0$ limit. The expression $\Psi(0, \dots, 0, w^k)$ means that we have evaluate the function $\Psi(w^1, \dots, w^k)$ at $w^1 = \dots = w^{k-1} = 0$ leaving it as a function only of w^k . In the original coordinates this is equivalent to setting $z^1 = \dots = z^{k-1} = z^k$.

Our goal is to show that $\epsilon \rightarrow 0$ limit of the right-hand side exists. The only ϵ dependence on the right hand side of (23) is in the integral over the regulation parameters t_1, \dots, t_k . Thus, it suffices to show that the $\epsilon \rightarrow 0$ limit of

$$\int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} \frac{dt_1 \dots dt_k}{(t_1 + \cdots + t_k)^d}$$

exists. By the AM/GM inequality we have $(t_1 + \cdots + t_k)^d \geq (t_1 \cdots t_k)^{d/k}$. So, the integral is bounded by

$$\int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} \frac{dt_1 \dots dt_k}{(t_1 + \cdots + t_k)^d} \leq \int_{(t_1, \dots, t_k) \in [\epsilon, L]^k} \frac{dt_1 \dots dt_k}{(t_1 \cdots t_k)^{d/k}} = \frac{1}{(1 - d/k)^k} \left(\epsilon^{1-d/k} - L^{1-d/k} \right)^k.$$

By assumption, $d < k$, so the right hand side has a well-defined $\epsilon \rightarrow 0$ limit. This concludes the proof. □

4. CHIRAL ANOMALIES IN ARBITRARY DIMENSIONS

Renormalization is an important step in constructing a quantum field theory. In the context of gauge theory, however, a consistent quantization requires that this renormalization behaves appropriately with respect to gauge symmetries present in the classical theory. This formalism for studying quantizations of gauge theories is due to Batalin-Vilkovisky [BV81], and has been made mathematically rigorous in the work of Costello

[Cos11]. The precise consistency of gauge symmetry with renormalization is encoded by the *quantum master equation*. Heuristically, one can think of the quantum master equation as a closedness condition on the path integral measure defined by the quantum action functional.

The key idea is the following: once a classical theory has been renormalized, so that we have a \hbar -linear effective family of functionals $\{I[L]\}$ whose $\hbar = 0$ limit is the classical action, the next step to constructing a quantization is to solve the quantum master equation (QME) for each functional $I[L]$. (In fact, once the QME holds at a single positive length $L > 0$ it holds for every other length by RG flow.) Often, the QME is not satisfied by the functional $I[L]$, but there exists a “correction” to $I[L]$ that does satisfy the QME. On the other hand, there may be unavoidable obstructions to solving this quantum master equation. These are known as *anomalies* in the physics literature. Since our method for solving the QME is deformation-theoretic in nature, these anomalies appear as cohomology classes in the cochain complex of local functionals.

In general, it is difficult to characterize such anomalies, but in the case of holomorphic theories on \mathbb{C}^d our result of one-loop finiteness from the previous section makes this problem much more tractable. Indeed, since there are no counterterms required, we can plug in the RG flow of the classical action functional and study the quantum master equation directly. As is usual in perturbation theory, one works order by order in \hbar to construct a quantization. However, in this section we continue to work linearly in \hbar , which is to say we study solutions to the quantum master equation modulo \hbar^2 .

4.1. The quantum master equation. In the BV formalism, as developed in [Cos11, CG17, CG], one has the following definition of a quantum field theory.

Definition 4.1. A *quantum field theory* in the BV formalism consists of a free BV theory (\mathcal{E}, Q, ω) and an effective family of functionals

$$\{I[L]\}_{L \in (0, \infty)} \subset \mathcal{O}_{P, sm}^+(\mathcal{E})[[\hbar]]$$

that satisfy:

(a) the exact renormalization group (RG) flow equation

$$I[L'] = W(P_{L < L'}, I[L]);$$

(b) the scale L quantum master equation (QME) at every length scale L :

$$(Q + \hbar \Delta_L) e^{I[L]/\hbar} = 0.$$

(c) as $L \rightarrow 0$, the functional $S[L]$ has an asymptotic expansion that is local.

The first part of the definition, namely RG flow, was the phenomena we studied in the previous section. We turn our attention to part two of the definition of a QFT. The

regularized quantum master equation at scale L can equivalently be written as

$$QI[L] + \hbar\Delta_L I[L] + \frac{1}{2}\{I[L], I[L]\}_L = 0;$$

Combined with part (c), the $\hbar \rightarrow 0, L \rightarrow 0$ limit of the above equation is precisely the classical master equation for the local functional $\lim_{L \rightarrow 0} I[L] \bmod \hbar$. A quantization of a classical functional $I \in \mathcal{O}_{\text{loc}}(\mathcal{E})$ is a quantization $\{I[L]\}$ as above whose $\hbar \rightarrow 0$ limit agrees with I .

In general, not every classical interaction admits a quantization. The *obstruction* to satisfying the quantum master equation order by order in \hbar is given by the following inductive definition.

Definition 4.2. Suppose $I[L] \in \mathcal{O}(\mathcal{E})[[\hbar]]/\hbar^{n+1}$ solves the QME modulo \hbar^n . The scale L obstruction to solving the QME modulo \hbar^n is

$$\Theta^{(n)}[L] = \hbar^{-n} \left(QI[L] + \hbar\Delta_L I[L] + \frac{1}{2}\{I[L], I[L]\}_L \right) \in \mathcal{O}(\mathcal{E}).$$

Equivalently, we can write the obstruction as $\Theta^{(n)}[L] = \hbar^{-n+1} e^{-I[L]/\hbar} (Q + \hbar\Delta_L) e^{I[L]/\hbar}$.

As a consequence of part (c) in the definition of a QFT above, the $L \rightarrow 0$ limit of the obstruction is defined and determines a cohomological degree +1 local functional

$$\Theta^{(n)} = \lim_{L \rightarrow 0} \Theta^{(n)}[L] \in \mathcal{O}_{\text{loc}}(\mathcal{E}).$$

Moreover, $\Theta^{(n)}$ is closed for the differential $Q + \{I, -\}$, where $I = \lim_{L \rightarrow 0} I[L] \bmod \hbar \in \mathcal{O}_{\text{loc}}(\mathcal{E})$.

In the remainder of this section we return to the holomorphic setting. Fix a classical holomorphic theory $(V, Q^{\text{hol}}, (-, -)_V, I^{\text{hol}})$ on \mathbb{C}^d . As usual, denote by $\mathcal{E}_V = (\Omega^{0,*}(\mathbb{C}^d, V), \bar{\partial} + Q^{\text{hol}})$ the linearized BRST complex of fields and $I = \int I^{\Omega^{0,*}}$ the classical interaction. Let $I[L] = \lim_{\epsilon \rightarrow 0} W(P_{\epsilon < L}, I) \bmod \hbar^2$ be the one-loop renormalization group flow using the propagator defined in Section 3.

4.2. The QME for holomorphic theories. The main result of this section is a characterization of the one-loop obstruction for holomorphic theories. Before jumping into the calculation, we state the following lemma, which is a simplification of the QME given our assumptions. Note that we only study one-loop effects here.

Lemma 4.3. *Let $\Theta[L] = \Theta^{(1)}[L]$ be the one-loop obstruction to the QME at scale L . Then, one has*

$$(27) \quad \hbar\Theta[L] = Q^{\text{hol}} I[L] + \frac{1}{2} \lim_{\epsilon \rightarrow 0} e^{-I/\hbar} e^{\hbar\partial_{P_{\epsilon < L}}} \left(\{I, I\}_{\epsilon} e^{I/\hbar} \right) \bmod \hbar^2.$$

Proof. We write the obstruction as $\Theta[L] = e^{-I[L]/\hbar} (Q + \hbar\Delta_L) e^{I[L]/\hbar}$. Notice that formally the one loop RG flow can equivalently be written as $e^{I[L]/\hbar} = \lim_{\epsilon \rightarrow 0} e^{\hbar\partial_{P_{\epsilon < L}}} e^{I/\hbar} \bmod \hbar^2$.

Applying the operator $Q + \hbar\Delta_L$ to both sides, we obtain

$$(Q + \hbar\Delta_L)e^{I[L]/\hbar} = \lim_{\epsilon \rightarrow 0} (Q + \hbar\Delta_L) \left(e^{\hbar\partial_{P_{\epsilon < L}}} e^{I/\hbar} \right).$$

The operator Q commutes with $e^{\hbar\partial_{P_{\epsilon < L}}}$. Moreover, one has $\Delta_L e^{\hbar\partial_{P_{\epsilon < L}}} = e^{\hbar\partial_{P_{\epsilon < L}}} \Delta_\epsilon$ as operators acting on functionals. Thus,

$$\lim_{\epsilon \rightarrow 0} (Q + \hbar\Delta_L) \left(e^{\hbar\partial_{P_{\epsilon < L}}} e^{I/\hbar} \right) = \lim_{\epsilon \rightarrow 0} e^{\hbar\partial_{P_{\epsilon < L}}} (Q + \hbar\Delta_\epsilon) e^{I/\hbar}.$$

Since Δ_ϵ is a BV operator with respect to the bracket $\{-, -\}_\epsilon$, we can rewrite the right-hand side as

$$\frac{1}{\hbar} \lim_{\epsilon \rightarrow 0} e^{\hbar P_{\epsilon < L}} (QI + \hbar\Delta_\epsilon I + \frac{1}{2}\{I, I\}_\epsilon) e^{I/\hbar}.$$

For every $\epsilon > 0$ we have $\Delta_\epsilon I = 0$. This is because I is a local functional and Δ_ϵ involves contraction with a factor of $\prod (d\bar{z}_i - d\bar{w}_i)$. Moreover, since I comes from a holomorphic Lagrangian we have $\bar{\partial}I = 0$.

Thus, the only terms remaining inside the parantheses in the above expression are $Q^{hol}I + \frac{1}{2}\{I, I\}_\epsilon$. We conclude that the obstruction $\Theta[L]$ can be expressed as

$$\begin{aligned} \Theta[L] &= \frac{1}{\hbar} \lim_{\epsilon \rightarrow 0} e^{-I/\hbar} e^{\hbar\partial_{P_{\epsilon < L}}} \left(Q^{hol}I + \frac{1}{2}\{I, I\}_\epsilon e^{I/\hbar} \right) \pmod{\hbar^2} \\ &= \frac{1}{\hbar} Q^{hol}I[L] + \frac{1}{2\hbar} \lim_{\epsilon \rightarrow 0} e^{-I/\hbar} e^{\hbar\partial_{P_{\epsilon < L}}} \left(\{I, I\}_\epsilon e^{I/\hbar} \right) \pmod{\hbar^2} \end{aligned}$$

as desired. In the second line, we have again used the fact that the operators Q^{hol} and $\partial_{P_{\epsilon < L}}$ commute. \square

As we saw above, the anomaly $\Theta[L]$ has a well-defined $L \rightarrow 0$ limit as a local functional and it is closed for the classical differential. Before stating the result, we need a modification of the definition of the weight of a given Feynman diagram. If Γ is a graph with a distinguished edge e , let $W_{\Gamma, e}(P_{\epsilon < L}, K_e, I)$ denote the weight of the graph as defined before, except with one minor difference. Instead of placing $P_{\epsilon < L}$ at each internal edge, we place K_e at the edge labeled e and $P_{\epsilon < L}$ on the remaining edges. The main result of this section is the following.

Proposition 4.4. *The obstruction $\Theta = \lim_{L \rightarrow 0} \Theta[L] \in \mathcal{O}_{loc}(\mathcal{E}_V)$ to satisfying the one-loop quantum master equation is given by the expression*

$$(28) \quad \hbar\Theta = Q^{hol} \lim_{L \rightarrow 0} I[L] + \frac{1}{2} \lim_{L \rightarrow 0} \lim_{\epsilon \rightarrow 0} \sum_{\Gamma \in \text{Wheel}_{d+1, e}} W_{\Gamma}(P_{\epsilon < L}, K_e, I)$$

where the sum is over all wheels with $(d+1)$ -vertices and distinguished edges thereof. In particular, when $Q^{hol} = 0$ (so that the first term vanishes), the anomaly is expressed as the sum over wheels with exactly $(d+1)$ -vertices.




FIGURE 1. The second term in Equation (28) representing the holomorphic anomaly.

This obstruction determines an element in the cohomology of the local deformation complex

$$[\Theta] \in H^1 \left(\mathcal{O}_{\text{loc}}(\mathcal{E}_V), \bar{\partial} + Q^{\text{hol}} + \{I, -\} \right).$$

This is a complete characterization of the cohomological obstruction to satisfying the quantum master equation for the classical theory I . If we chose any other quantization of $\{I'[L]\}$ of I , say coming from a different gauge fixing condition, we obtain class cohomologous to this $[\Theta] = [\Theta']$.

Proof of Proposition 4.4. Like the proof of the non-existence of counterterms for holomorphic theories, the proof of this result will be the consequence of an explicit calculations and bounds of certain Feynman diagrams.

Note that the first term, involving Q^{hol} , is the $L \rightarrow 0$ limit of the right-hand side of Equation (27). Thus, it suffices to focus on the second term.

We express the quantity

$$(29) \quad \lim_{\epsilon \rightarrow 0} e^{-I/\hbar} e^{\hbar \partial_{P_{\epsilon < L}}} \left(\{I, I\}_{\epsilon} e^{I/\hbar} \right) \quad \text{mod } \hbar^2$$

as a sum over graphs. By assumption, we are only looking at graphs of genus one which look like wheels with possible trees attach. Graphically, the quantity $\{I, I\}_{\epsilon}$ is the graph of two vertices with a separating edge labeled by the heat kernel K_{ϵ} . Thus, all weights appearing in the expansion of (29) attach the propagator $P_{\epsilon < L}$ to all edges besides a single distinguished edge e , which is labeled by K_{ϵ} . Thus, as a over a sum of graphs, we see that the following two types of weights occur in the expansion of (29).

- (a) the distinguished edge e is separating;
- (b) the distinguished edge e is *not* separating, and so appears as the internal edge of the wheel portion of the graph.

By the classical master equation, we see that the $\epsilon \rightarrow 0$ limit of weights of Type (a) go to zero. Thus, we must only consider the weights of Type (b).

The result will follow from two steps. These should seem familiar from the proof of the main result about the existence of no counterterms.

- (1) If Γ is a wheel with $k < d + 1$ vertices, then $W_\Gamma(P_{\epsilon < L}, K_\epsilon, I) = 0$ identically.
- (2) If Γ is a wheel with $k > d + 1$ vertices, then $\lim_{\epsilon \rightarrow 0} W_\Gamma(P_{\epsilon < L}, K_\epsilon, I) = 0$.

The proof of both of these facts is only dependent on the analytic part of the weights. Thus, it suffices to make the same reduction as we did in the previous section. To extract that analytic part of the graph we proceed as in Definition 3.10. If $(k, (\vec{n}))$ is a pair of wheel data (recall k labels the number of vertices and \vec{n} labels the derivatives at each vertex) define the smooth distribution

$$\tilde{W}_{\epsilon < L}^{k, (n)} : C_c^\infty((\mathbb{C}^d)^k) \rightarrow \mathbb{C},$$

that sends a smooth compactly supported function $\Phi \in C_c^\infty((\mathbb{C}^d)^k) = C_c^\infty(\mathbb{C}^{dk})$ to

$$(30) \quad \tilde{W}_{\epsilon < L}^{k, (n)}(\Phi) = \int_{(z^1, \dots, z^k) \in (\mathbb{C}^d)^k} \prod_{\alpha=1}^k d^d z^\alpha \Phi(z^1, \dots, z^\alpha) \left(\frac{\partial}{\partial z^k} \right)^{\vec{n}^k} K_\epsilon(z^1, z^k) \prod_{\alpha=1}^{k-1} \left(\frac{\partial}{\partial z^\alpha} \right)^{\vec{n}^\alpha} P_{\epsilon < L}^{an}(z^\alpha, z^{\alpha+1}).$$

Item (1) follows from the following observation.

Lemma 4.5. *Let $(k, (\vec{n}))$ be a pair of wheel data. If the number of vertices k satisfies $k \leq d$ then*

$$\tilde{W}_{\epsilon < L}^{k, (n)} = 0$$

as a distribution on \mathbb{C}^{dk} for any $\epsilon, L > 0$.

Proof. In fact, the integrand of (30) is identically zero provided $k \leq d$ by a simple observation of the differential form type. Consider the factor in the integrand of $\tilde{W}_{\epsilon < L}^{k, (n)}$ given by

$$\left(\frac{\partial}{\partial z^k} \right)^{\vec{n}^k} K_\epsilon(z^1, z^k) \prod_{\alpha=1}^{k-1} \left(\frac{\partial}{\partial z^\alpha} \right)^{\vec{n}^\alpha} P_{\epsilon < L}^{an}(z^\alpha, z^{\alpha+1}).$$

Making the usual change of coordinates $w^\alpha = z^{\alpha+1} - z^\alpha$ and $w^k = z^k$ we see that this factor is proportional to the following constant coefficient differential form

$$\left(\prod_{i=1}^d \left(\sum_{\alpha=1}^{k-1} d\bar{w}_i^\alpha \right) \right) \prod_{\alpha=1}^{k-1} \left(\eta^\alpha \prod_{i=1}^d d\bar{w}_i^\alpha \right).$$

Note that this differential form only involves the coordinates (w_i^α) for $\alpha = 1, \dots, k-1$. Thus, we may consider it as a Dolbeault form on $\mathbb{C}^{d(k-1)}$. As such, it is of the type $(0, d + (k-1)(d-1)) = (0, (d-k+1) + d(k-1))$. Clearly, $(d-k+1) + d(k-1) > d(k-1)$ provided $k \leq d$. Thus, the weight is identically zero provided $k \leq d$, as desired. \square

Item (2) follows from the following technical lemma that the analytic weight associated to the wheels of valency $k > d + 1$ vanish in the limit $\epsilon \rightarrow 0$.

Lemma 4.6. Let $(k, (\vec{n}))$ be a pair of wheel data such that $k > d + 1$. Then the $\epsilon \rightarrow 0$ limit of the analytic weight

$$\lim_{\epsilon \rightarrow 0} \tilde{W}_{\epsilon < L}^{k, (n)} = 0$$

is identically zero as a distribution on \mathbb{C}^{dk} .

Proof. The proof is very similar to the argument we gave in the proof of Lemma 3.12, so we will be a bit more concise. First, we make the familiar change of coordinates as in Equations (18),(19). Using the explicit form the heat kernel and propagator we see that for any $\Phi \in C_c^\infty(\mathbb{C}^{dk})$ the weight is

$$\begin{aligned} \tilde{W}_{\epsilon < L}^{k, (n)}(\Phi) &= \int_{w^k \in \mathbb{C}^d} d^{2d} w^k \int_{(w_1, \dots, w_{k-1}) \in (\mathbb{C}^d)^{k-1}} \left(\prod_{\alpha=1}^{k-1} d^{2d} w^\alpha \right) \Phi(w^1, \dots, w^k) \\ &\times \int_{(t_1, \dots, t_k) \in [\epsilon, L]^{k-1}} \frac{1}{(2\pi i \epsilon)^d} \prod_{\alpha=1}^{k-1} \frac{dt_\alpha}{(2\pi i t_\alpha)^d} \\ &\times \sum_{i_1, \dots, i_{k-1}=1}^d \epsilon_{i_1, \dots, i_d} \left(\frac{\bar{w}_1^1}{t_1} \frac{(\bar{w}^1)^{n^1}}{4t^{n^1}} \right) \cdots \left(\frac{\bar{w}_{k-1}^{k-1}}{4t_{k-1}} \frac{(\bar{w}^{k-1})^{n^{k-1}}}{4t^{n^{k-1}}} \right) \left(\frac{1}{4t^{n^k}} \left(\sum_{\alpha=1}^{k-1} \bar{w}^\alpha \right)^{n^k} \right) \\ &\times \exp \left(- \sum_{\alpha=1}^{k-1} \frac{|w^\alpha|^2}{4t_\alpha} - \frac{1}{4\epsilon} \left| \sum_{\alpha=1}^{k-1} w^\alpha \right|^2 \right). \end{aligned}$$

We will integrate by parts to eliminate the factors of \bar{w}_i^α .

For each $1 \leq \alpha < k$ and i_α , define the ϵ and $t = (t_1, \dots, t_{k-1})$ -dependent holomorphic differential operator

$$D_{\alpha, i_\alpha}(t) := \left(\frac{\partial}{\partial w_{i_\alpha}^\alpha} - \sum_{\beta=1}^{k-1} \frac{t_\beta}{t_1 + \cdots + t_{k-1} + \epsilon} \frac{\partial}{\partial w_{i_\alpha}^\beta} \right) \prod_{j=1}^d \left(\frac{\partial}{\partial w_j^\alpha} - \sum_{\beta=1}^{k-1} \frac{t_\beta}{t_1 + \cdots + t_{k-1} + \epsilon} \frac{\partial}{\partial w_j^\beta} \right)^{n_j^\alpha}.$$

And the ϵ, t -dependent holomorphic differential operator

$$D_k(t) = \prod_{j=1}^d \left(\frac{\partial}{\partial w_j^k} - \sum_{\beta=1}^{k-1} \frac{t_\beta}{t_1 + \cdots + t_{k-1} + \epsilon} \frac{\partial}{\partial w_j^\beta} \right)^{n_j^k}.$$

By a completely analogous version of Lemma the operators above allow us to integrate by parts and express the weight in the form

$$\begin{aligned} \tilde{W}_{\epsilon < L}^{k, (n)}(\Phi) &= \pm \int_{w^k \in \mathbb{C}^d} d^{2d} w^k \int_{(w_1, \dots, w_{k-1}) \in (\mathbb{C}^d)^{k-1}} \left(\prod_{\alpha=1}^{k-1} d^{2d} w^\alpha \right) \\ &\times \int_{(t_1, \dots, t_{k-1}) \in [\epsilon, L]^{k-1}} \frac{1}{(2\pi i \epsilon)^d} \prod_{\alpha=1}^{k-1} \frac{dt_\alpha}{(2\pi i t_\alpha)^d} \\ &\times \left(\sum_{i_1, \dots, i_{k-1}} \epsilon_{i_1, \dots, i_d} D_{1, i_1}(t) \cdots D_{k-1, i_{k-1}}(t) D_k(t) \Phi(w^1, \dots, w^k) \right) \\ &\times \exp \left(- \sum_{\alpha=1}^{k-1} \frac{|w^\alpha|^2}{4t_\alpha} - \frac{1}{4\epsilon} \left| \sum_{\alpha=1}^{k-1} w^\alpha \right|^2 \right). \end{aligned}$$

Observe that the operators $D_{\alpha, i_\alpha}(t), D_k(t)$ are uniformly bounded in t . Thus, there exists a constant $C = C(\Phi) > 0$ depending only on the function Φ such that we can bound the weight as

$$(31) \quad |\widetilde{W}_{\epsilon < L}^{k, (n)}(\Phi)| \leq C \int_{(w^1, \dots, w^{k-1})} \prod_{\alpha=1}^{k-1} d^{2d} w^\alpha \int_{(t_1, \dots, t_{k-1}) \in [\epsilon, L]^{k-1}} dt_1 \dots dt_k \frac{1}{\epsilon^d t_1^d \dots t_{k-1}^d} \\ \times \exp \left(- \sum_{\alpha=1}^{k-1} \frac{|w^\alpha|^2}{4t_\alpha} - \frac{1}{4\epsilon} \left| \sum_{\alpha=1}^{k-1} w^\alpha \right|^2 \right).$$

Thus, to show that the limit $\lim_{L \rightarrow 0} \lim_{\epsilon \rightarrow 0} \widetilde{W}_{\epsilon < L}^{k, (n)}(\Phi) = 0$ it suffices to show that the limit of the right-hand side vanishes.

The Gaussian integral over the variables w_i^α contributes the following factor

$$\int_{(w^1, \dots, w^{k-1})} \prod_{\alpha=1}^{k-1} d^{2d} w^\alpha \exp \left(- \sum_{\alpha=1}^{k-1} \frac{|w^\alpha|^2}{4t_\alpha} - \frac{1}{4\epsilon} \left| \sum_{\alpha=1}^{k-1} w^\alpha \right|^2 \right) = C' \left(\frac{\epsilon t_1 \dots t_{k-1}}{\epsilon + t_1 + \dots + t_{k-1}} \right)^d.$$

Where C' involves factors of 2 and π . Plugging this back in to the right-hand side of (31) we see that

$$|\widetilde{W}_{\epsilon < L}^{k, (n)}(\Phi)| \leq CC' \int_{[\epsilon, L]^{k-1}} \frac{dt_1 \dots dt_{k-1}}{(\epsilon + t_1 + \dots + t_{k-1})^d} \leq CC' \prod_{\alpha=1}^{k-1} \int_{t_\alpha=\epsilon}^L dt_\alpha t_\alpha^{-d/(k-1)}.$$

In the second inequality we have used the fact that $\epsilon > 0$ and the AM-GM inequality. It is immediate to see that the $\epsilon \rightarrow 0$ limit of the above exists provided $k > d + 1$, which is the situation we are in, and that the $L \rightarrow 0$ limit vanishes. \square

This completes the proof of Proposition 4.4 \square

4.2.1. *Relation to the ABJ anomaly.* The lemma we have just proved implies that for holomorphic theories on \mathbb{C}^d the anomaly is given by evaluating a collection of wheel diagrams with exactly $d + 1$ vertices. This expression for the obstruction fits into a generic class of one-loop anomalies from gauge theory called the Adler-Bell-Jackiw (ABJ) anomaly [Adl69, BJ69]. This anomaly is most commonly associated with four dimensional gauge theory.

We recall the basic setup for the ABJ anomaly. Consider a free Dirac fermion Ψ on \mathbb{R}^4 coupled to a background gauge field $A \in \Omega^1(\mathbb{R}^4) \otimes \mathfrak{g}$. For this to make sense, Ψ is taken to be in valued in a representation V of the Lie algebra \mathfrak{g} so we may think of it as an element $\Psi \in \mathcal{S}(\mathbb{R}^4) \otimes V$. Here, $\mathcal{S}(\mathbb{R}^4)$ is the space of sections of the full spinor bundle on \mathbb{R}^4 . The action functional is

$$S(A, \Psi) = \int \langle \Psi, \not{\partial}_A \Psi \rangle_V$$

where $\not{\partial}_A = \not{\partial} + [A, -]$ is the A -coupled Dirac operator. We are implicitly using the canonical spin invariant symplectic pairing $\mathcal{S} \otimes \mathcal{S} \rightarrow \Omega^4(\mathbb{R}^4) = C^\infty(\mathbb{R}^4) d^4x$ and a \mathfrak{g} -invariant pairing $\langle -, - \rangle_V : V \otimes V \rightarrow \mathbb{C}$, to obtain a local functional.

For any smooth map $\alpha : \mathbb{R}^4 \rightarrow \mathfrak{g}$, the infinitesimal transformation $\Psi \rightarrow \Psi + \epsilon[\alpha, \Psi]$ (where ϵ is an even parameter of square zero) is a classical symmetry of $S(A, \Psi)$. Quantum mechanically, there is a one-loop anomaly which measures the failure of the path integral to be invariant with respect to this symmetry. It is a well-known calculation, see for instance [FS04], that this anomaly is measured by the following local functional

$$(32) \quad \int \text{Tr}_V(\alpha F_A F_A).$$

The trace is taken in the representation V . The fundamental calculation is the infamous “triangle diagram”, where two vertices are labeled by the gauge field and the third by α . In practice, physicists express the anomaly as a failure for the Noether current associated to the symmetry α to be divergenceless.

There is the following holomorphic version of this anomaly. Again, let V be a \mathfrak{g} representation. Consider the following action functional on \mathbb{C}^2 :

$$S(A, \beta, \gamma) = \int \langle \beta, \bar{\partial}_A \gamma \rangle_V$$

where $\gamma : \mathbb{C}^2 \rightarrow V$, $\beta \in \Omega^{2,1}(\mathbb{C}^2, V)$, and $A \in \Omega^{0,1}(\mathbb{C}^2, \mathfrak{g})$. Since A is a $(0,1)$ form it defines a deformation of the trivial holomorphic G -bundle. Although we have not put this theory in the BV formalism, there is a natural way to do so. The infinitesimal symmetry we contemplate is of the form $\gamma \rightarrow \gamma + \epsilon[\alpha, \gamma]$ where $\alpha : \mathbb{C}^2 \rightarrow \mathfrak{g}$. We study the anomaly to quantizing this symmetry to one-loop. Following the result for the anomaly given in the previous section, one sees that it is computed by a wheel with three vertices. For type reasons, one vertex is labeled α and the other two are labeled by the gauge fields A .

$$\int \text{Tr}_V(\alpha \partial A \partial A).$$

This is the holomorphic version of ABJ anomaly (32). Note that there are no terms of order A^3 or above. In fact, the functional $\int \text{Tr}(\alpha F_A F_A)$ is cohomologous to the expression above in the local deformation complex.

Remark 4.7. We have already shown how familiar topological theories can be cast in a holomorphic language. For instance, topological BF theory is a holomorphic deformation of holomorphic BF theory. It is a peculiar consequence of the above result that such topological theories also admit a simple regularization procedure. Without much more difficulty, one can extend this to certain topological theories to odd dimensional manifolds of the form $X \times S$, where X is a complex manifold and S is a real one-dimensional manifold. We consider the theory as a product of a holomorphic theory on X and a one-dimensional topological theory on S . This can be further extended to transversely holomorphic foliations [Bru96, Ghy96], which we will study in a future publication. Further, often topological BF theory further deforms to Yang-Mills. It would be interesting to apply our analysis above to such gauge theories.

APPENDIX A. SOME FUNCTIONAL ANALYSIS

Homological algebra plays a paramount role in our approach to quantum field theory. We immediately run into a subtle issue, which is that the underlying graded spaces of the complexes of fields we are interested in are infinite dimensional, so care must be taken when defining constructions such as duals and homomorphism spaces. A common approach to dealing with issues of infinite dimensional linear algebra is to consider vector spaces equipped with a topology. A problem with this is that the category of topological vector spaces is not an abelian category, so doing any homological algebra in this naive category is utterly hopeless. It is therefore advantageous to enlarge this to the category of *differentiable vector spaces*. The details of this setup are carried out in the Appendix of [CG17], but we will recall some key points for completeness of exposition. In this appendix we also set up our notation for duals and function spaces.

Let \mathbf{Mfld} be the site of smooth manifolds. The covers defining the Grothendieck topology are given by surjective local diffeomorphisms. There is a natural sheaf of algebras on this site given by smooth functions $C^\infty : M \mapsto C^\infty(M)$.

For any p the assignment $\Omega^p : M \mapsto \Omega^p(M)$ defines a C^∞ -module. Similarly, if F is any C^∞ -module we have the C^∞ -module of p -forms with values on F defined by the assignment

$$\Omega^p(F) : M \in \mathbf{Mfld} \mapsto \Omega^p(M, F) = \Omega^p(M) \otimes_{C^\infty(M)} F(M).$$

Definition A.1. A *differentiable vector space* is a C^∞ -module equipped with a map of sheaves on \mathbf{Mfld}

$$\nabla : F \rightarrow \Omega^1(F)$$

such that for each M , $\nabla(M)$ defines a flat connection on the $C^\infty(M)$ -module $F(M)$. A map of differentiable vector spaces is one of C^∞ -modules that intertwines the flat connections. This defines a category that we denote \mathbf{DVS} .

Our favorite example of differentiable vector spaces are imported directly from geometry.

Example A.2. Suppose E is a vector bundle on a manifold X . Let $\mathcal{E}(X)$ denote the space of smooth global sections. Let $C^\infty(M, \mathcal{E}(X))$ be the space of sections of the bundle $\pi_X^* E$ on $M \times X$ where $\pi_X : M \times X \rightarrow X$ is projection. The assignment $M \mapsto C^\infty(M, \mathcal{E}(X))$ is a C^∞ -module with flat connection, so defines a differentiable vector space. Similarly, the space of compactly supported sections $\mathcal{E}_c(X)$ is a \mathbf{DVS} .

Many familiar categories of topological vector spaces embed inside the category of differentiable vector spaces. Consider the category of locally convex topological vector spaces \mathbf{LCTVS} . If V is such a vector space, there is a notion of a smooth map $f : U \subset \mathbb{R}^n \rightarrow V$. One can show, Proposition B.3.0.6 of [CG17], that this defines a functor $\mathbf{dif}_f : \mathbf{LCTVS} \rightarrow \mathbf{DVS}$.

LCTVS \rightarrow DVS sending V to the C^∞ -module $M \mapsto C^\infty(M, V)$. If $BVS \subset LCTVS$ is the subcategory with the same objects but whose morphisms are bounded linear maps, this functor restricts to embed BVS as a full subcategory $BVS \subset DVS$.

There is a notion of completeness that is useful when discussing tensor products. A topological vector space $V \in BVS$ is *complete* if every smooth map $c : \mathbb{R} \rightarrow V$ has an anti-derivative [KM97]. There is a full subcategory $CVS \subset BVS$ of complete topological vector spaces. The most familiar example of a complete topological vector space will be the smooth sections $\mathcal{E}(X)$ of a vector bundle $E \rightarrow X$.

We let $\text{Ch}(DVS)$ denote the category of cochain complexes in differentiable vector spaces (we will refer to objects as differentiable vector spaces). It is enriched over the category of differential graded vector spaces in the usual way. We say that a map of differentiable cochain complexes $f : V \rightarrow W$ is a quasi-isomorphism if and only if for each M the map $f : C^\infty(M, V) \rightarrow C^\infty(M, W)$ is a quasi-isomorphism.

Theorem A.3 (Appendix B [CG17]). *The full subcategory $\text{dif}_c : CVS \subset DVS$ is closed under limits, countable coproducts, and sequential colimits of closed embeddings. Furthermore, CVS has the structure of a symmetric monoidal category with respect to the completed tensor product $\widehat{\otimes}_\beta$.*

We will not define the tensor product $\widehat{\otimes}_\beta$ here, but refer the reader the cited reference for a complete exposition. We will recall its key properties below. Often times we will write \otimes for $\widehat{\otimes}_\beta$ where there is no potential conflict of notation. The fundamental property of the tensor product that we use is the following. Suppose that E, F are vector bundles on manifolds X, Y respectively. Then, $\mathcal{E}(X), \mathcal{F}(Y)$ lie in CVS , so it makes sense to take their tensor product using $\widehat{\otimes}_\beta$. There is an isomorphism

$$(33) \quad \mathcal{E}(X) \widehat{\otimes}_\beta \mathcal{F}(Y) \cong \Gamma(X \times Y, E \boxtimes F)$$

where $E \boxtimes F$ denotes the external product of bundles, and Γ is smooth sections.

If E is a vector bundle on a manifold X , then the spaces $\mathcal{E}(X), \mathcal{E}_c(X)$ both lie in the subcategory $CVS \subset DVS$. The differentiable structure arises from the natural topologies on the spaces of sections.

We will denote by $\overline{\mathcal{E}}(X)$ ($\overline{\mathcal{E}}_c(X)$) the space of (compactly supported) distributional sections. It is useful to bear in mind the following inclusions

$$\begin{array}{ccc}
 & \overline{\mathcal{E}}_c(X) & \\
 \nearrow & & \searrow \\
 \mathcal{E}_c(X) & & \overline{\mathcal{E}}(X) \\
 \searrow & & \nearrow \\
 & \mathcal{E}(X) & .
 \end{array}$$

When X is compact the bottom left and top right arrows are equalities.

Denote by E^\vee the dual vector bundle whose fiber over $x \in X$ is the linear dual of E_x . Let $E^!$ denote the vector bundle $E^\vee \otimes \text{Dens}_X$, where Dens_X is the bundle of densities. In the case X is oriented, Dens_X is isomorphic to the top wedge power of T^*X . Let $\mathcal{E}^!(X)$ denote the space of sections of $E^!$. The natural pairing

$$\mathcal{E}_c(X) \otimes \mathcal{E}^!(X) \rightarrow \mathbb{C}$$

that pairs sections of E with the evaluation pairing and integrates the resulting compactly supported top form exhibits $\bar{\mathcal{E}}_c(X)$ as the continuous dual to $\mathcal{E}^!(X)$. Likewise, $\mathcal{E}_c(X)$ is the continuous dual to $\bar{\mathcal{E}}^!(X)$. In this way, the topological vector spaces $\bar{\mathcal{E}}(X)$ and $\bar{\mathcal{E}}_c(X)$ obtain a natural differentiable structure.

If V is any differentiable vector space then we define the space of linear functionals on V to be the space of maps $V^* = \text{Hom}_{\text{DVS}}(V, \mathbb{R})$. Since DVS is enriched over itself this is again a differentiable vector space. Similarly, we can define the polynomial functions of homogeneous degree n to be the space

$$\text{Sym}^n(V^*) = \text{Hom}_{\text{DVS}}^{\text{multi}}(V \times \cdots \times V, \mathbb{R})_{S_n}$$

where the hom-space denotes multi-linear maps, and we have taken S_n -coinvariants on the right-hand side. The algebra of functions on V is defined by

$$\mathcal{O}(V) = \prod_n \text{Sym}^n(V^*).$$

As an application of Equation (33) we have the following identification.

Lemma A.4. *Let E be a vector bundle on X . Then, there is an isomorphism*

$$\mathcal{O}(\mathcal{E}(X)) \cong \prod_n \mathcal{D}_c(X^n, (E^!)^{\boxtimes n})_{S_n}$$

where $\mathcal{D}_c(X^n, (E^!)^{\boxtimes n})$ is the space of compactly supported distributional sections of the vector bundle $(E^!)^{\boxtimes n}$. Again, we take S_n -coinvariants on the right hand side.

REFERENCES

- [Adl69] Stephen L. Adler. Axial vector vertex in spinor electrodynamics. *Phys. Rev.*, 177:2426–2438, 1969.
- [BCOV94] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa. Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes. *Comm. Math. Phys.*, 165(2):311–427, 1994.
- [BD04] Alexander Beilinson and Vladimir Drinfeld. *Chiral algebras*, volume 51 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
- [BJ69] J. S. Bell and R. Jackiw. A PCAC puzzle: $\pi \rightarrow \gamma\gamma$ in the sigma model. *Nuovo Cim.*, A60:47–61, 1969.
- [BR73] I. N. Bernstein and B. I. Rosenfeld. Homogeneous spaces of infinite-dimensional Lie algebras and the characteristic classes of foliations. *Uspehi Mat. Nauk*, 28(4(172)):103–138, 1973.
- [Bru96] Marco Brunella. On transversely holomorphic flows. I. *Invent. Math.*, 126(2):265–279, 1996.
- [BV81] I. A. Batalin and G. A. Vilkovisky. Gauge algebra and quantization. *Phys. Lett. B*, 102(1):27–31, 1981.

- [CG] Kevin Costello and Owen Gwilliam. Factorization algebras in quantum field theory. Vol. 2. available at <http://people.mpim-bonn.mpg.de/gwilliam>.
- [CG17] Kevin Costello and Owen Gwilliam. *Factorization algebras in quantum field theory. Vol. 1*, volume 31 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2017.
- [CL] Kevin Costello and Si Li. Twisted supergravity and its quantization. Available at <https://arxiv.org/abs/1606.00365>.
- [Cosa] Kevin Costello. Holography and Koszul duality: the example of the M2 brane. Available at <https://arxiv.org/abs/1705.02500>.
- [Cosb] Kevin Costello. M-theory in the Omega-background and 5-dimensional non-commutative gauge theory. Available at <https://arxiv.org/abs/1610.04144>.
- [Cosc] Kevin Costello. Supersymmetric gauge theory and the Yangian. Available at <http://arxiv.org/abs/1303.2632>.
- [Cos11] Kevin Costello. *Renormalization and effective field theory*, volume 170 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2011.
- [Cos13] Kevin Costello. Notes on supersymmetric and holomorphic field theories in dimensions 2 and 4. *Pure Appl. Math. Q.*, 9(1):73–165, 2013.
- [CW04] Karen Chandler and Pit-Mann Wong. Finsler geometry of holomorphic jet bundles. In *A sampler of Riemann-Finsler geometry*, volume 50 of *Math. Sci. Res. Inst. Publ.*, pages 107–196. Cambridge Univ. Press, Cambridge, 2004.
- [ES] Richard Eager and Ingmar Saberi. Holomorphic field theories and Calabi–Yau algebras. Available at <https://arxiv.org/abs/1805.02084>.
- [EWY18] Chris Elliott, Brian Williams, and Philsang Yoo. Asymptotic freedom in the BV formalism. *J. Geom. Phys.*, 123:246–283, 2018.
- [FG12] John Francis and Dennis Gaitsgory. Chiral Koszul duality. *Selecta Math. (N.S.)*, 18(1):27–87, 2012.
- [FS04] Kazuo Fujikawa and Hiroshi Suzuki. *Path integrals and quantum anomalies*, volume 122 of *International Series of Monographs on Physics*. The Clarendon Press, Oxford University Press, New York, 2004. Translated from the 2001 Japanese original.
- [Fuk86] D. B. Fuks. *Cohomology of infinite-dimensional Lie algebras*. Contemporary Soviet Mathematics. Consultants Bureau, New York, 1986. Translated from the Russian by A. B. Sosinskiĭ.
- [Get09] Ezra Getzler. Lie theory for nilpotent L_∞ -algebras. *Ann. of Math. (2)*, 170(1):271–301, 2009.
- [GG80] Mark Green and Phillip Griffiths. Two applications of algebraic geometry to entire holomorphic mappings. In *The Chern Symposium 1979 (Proc. Internat. Sympos., Berkeley, Calif., 1979)*, pages 41–74. Springer, New York-Berlin, 1980.
- [GGW] Owen Gwilliam, Vassily Gorbounov, and Brian Williams. Chiral differential operators via Batalin-Vilkovisky quantization. Available at <https://arxiv.org/abs/1610.09657>.
- [GH94] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994. Reprint of the 1978 original.
- [Ghy96] Étienne Ghys. On transversely holomorphic flows. II. *Invent. Math.*, 126(2):281–286, 1996.
- [Gui73] Victor W. Guillemin. Cohomology of vector fields on a manifold. *Advances in Math.*, 10:192–220, 1973.
- [Joh94] A. Johansen. Twisting of N=1 susy gauge theories and heterotic topological theories. arXiv:hep-th/9403017, 1994.
- [KM97] Andreas Kriegl and Peter W. Michor. *The convenient setting of global analysis*, volume 53 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [Li] Si Li. Vertex algebras and quantum master equation. available at <https://arxiv.org/abs/1612.01292>.

- [LMNS96] Andrei Losev, Gregory Moore, Nikita Nekrasov, and Samson Shatashvili. Four-dimensional avatars of two-dimensional RCFT. *Nuclear Phys. B Proc. Suppl.*, 46:130–145, 1996. S-duality and mirror symmetry (Trieste, 1995).
- [LMNS97] Andrei Losev, Gregory Moore, Nikita Nekrasov, and Samson Shatashvili. Chiral Lagrangians, anomalies, supersymmetry, and holomorphy. *Nuclear Phys. B*, 484(1-2):196–222, 1997.
- [Los98] Mark V. Losik. Diagonal cohomology of the Lie algebra of vector fields. *Differential Geom. Appl.*, 8(1):71–86, 1998.
- [Nek96] Nikita Alex Nekrasov. *Four-dimensional holomorphic theories*. ProQuest LLC, Ann Arbor, MI, 1996. Thesis (Ph.D.)—Princeton University.
- [Pol98] Joseph Polchinski. *String theory. Vol. I*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1998. An introduction to the bosonic string.
- [Pop99] A. D. Popov. Holomorphic Chern-Simons-Witten theory: from 2D to 4D conformal field theories. *Nuclear Phys. B*, 550(3):585–621, 1999.
- [Pop00] Alexander D. Popov. Holomorphic analogs of topological gauge theories. *Phys. Lett. B*, 473(1-2):65–72, 2000.
- [Sta92] Jim Stasheff. Differential graded Lie algebras, quasi-Hopf algebras and higher homotopy algebras. In *Quantum groups (Leningrad, 1990)*, volume 1510 of *Lecture Notes in Math.*, pages 120–137. Springer, Berlin, 1992.
- [tHV74] G. 't Hooft and M. Veltman. One-loop divergencies in the theory of gravitation. *Ann. Inst. H. Poincaré Sect. A (N.S.)*, 20:69–94, 1974.
- [Wil17] Brian Williams. The Virasoro vertex algebra and factorization algebras on Riemann surfaces. *Letters in Mathematical Physics*, Aug 2017. <https://doi.org/10.1007/s11005-017-0982-7>.
- [Wit88] Edward Witten. Topological quantum field theory. *Comm. Math. Phys.*, 117(3):353–386, 1988.

NORTHEASTERN UNIVERSITY

Email address: brwilliams@northeastern.edu